

| Indicators | Chhattisgarh | | Koriya | | Surguja | | Jashpur | |
|--|--------------|-------|--------|-------|---------|-------|---------|-------|
| | Total | Rural | Total | Rural | Total | Rural | Total | Rural |
| Nutritional status below 5 years | | | | | | | | |
| Below -2 SD Wasting (Weight for Height) (%) | | | | | | | | |
| Male | 31.8 | 30.8 | 13.6 | 13.3 | 21.6 | 21.2 | 25.0 | 25.2 |
| Female | 33.1 | 32.7 | 14.1 | 13.7 | 24.8 | 24.7 | 26.5 | 26.6 |
| Person | 32.4 | 31.7 | 13.8 | 13.6 | 23.0 | 22.7 | 25.7 | 25.9 |
| Below -3 SD Wasting (Weight for Height) (%) | | | | | | | | |
| Male | 11.8 | 11.2 | 4.2 | 4.5 | 2.0 | 1.9 | 1.9 | 1.9 |
| Female | 11.1 | 10.9 | - | - | 3.2 | 3.4 | 4.3 | 4.4 |
| Person | 11.5 | 11.1 | 2.7 | 2.7 | 2.5 | 2.5 | 3.0 | 3.1 |
| Below -2 SD Stunting (Height for Age) (%) | | | | | | | | |
| Male | 35.9 | 36.8 | 43.8 | 42.7 | 27.0 | 27.6 | 25.7 | 25.8 |
| Female | 33.4 | 34.2 | 47.9 | 47.0 | 20.7 | 21.0 | 17.9 | 17.7 |
| Person | 34.7 | 35.5 | 45.7 | 44.7 | 24.2 | 24.7 | 22.1 | 22.1 |
| Below -3SD Stunting (Height for Age) (%) | | | | | | | | |
| Male | 21.2 | 21.5 | 23.4 | 22.2 | 14.9 | 15.5 | 9.7 | 9.8 |
| Female | 18.7 | 18.2 | 26.5 | 24.0 | 9.1 | 9.4 | 4.2 | 3.9 |
| Person | 20.0 | 19.9 | 24.8 | 23.0 | 12.3 | 12.8 | 7.2 | 7.1 |
| Below -2 SD Underweight (Weight for Age) (%) | | | | | | | | |
| Male | 41.3 | 41.8 | 38.4 | 37.5 | 24.7 | 24.3 | 26.5 | 26.6 |
| Female | 37.1 | 37.3 | 43.7 | 41.5 | 19.1 | 19.6 | 19.0 | 18.9 |
| Person | 39.4 | 39.7 | 40.8 | 39.3 | 22.2 | 22.2 | 23.0 | 23.1 |
| Below -3 SD Underweight (Weight for Age) (%) | | | | | | | | |
| Male | 19.9 | 19.1 | 15.2 | 14.0 | 8.3 | 8.5 | 5.4 | 5.5 |
| Female | 17.4 | 17.0 | 17.8 | 14.4 | 5.9 | 6.1 | - | - |
| Person | 18.8 | 18.1 | 16.4 | 14.2 | 7.2 | 7.4 | 4.0 | 4.1 |
| Below -2 SD Undernourished (BMI for Age) (%) | | | | | | | | |
| Male | 32.7 | 31.4 | 10.2 | 10.1 | 21.8 | 21.6 | 27.5 | 27.8 |
| Female | 34.2 | 33.4 | 13.8 | 13.5 | 26.5 | 26.2 | 33.4 | 33.7 |
| Person | 33.5 | 32.4 | 11.8 | 11.6 | 23.9 | 23.6 | 30.2 | 30.5 |
| Below -3 SD Undernourished (BMI for Age) (%) | | | | | | | | |
| Male | 16.5 | 14.9 | 5.5 | 5.9 | 2.6 | 2.4 | - | - |
| Female | 15.1 | 14.0 | 3.2 | - | 5.0 | 5.2 | 7.1 | 7.1 |
| Person | 15.9 | 14.5 | 4.5 | 4.7 | 3.7 | 3.6 | 5.3 | 5.3 |
| Above 2 SD Overnourished (BMI for Age) (%) | | | | | | | | |
| Male | 7.9 | 8.4 | 9.1 | 9.9 | 8.5 | 8.9 | - | - |
| Female | 7.4 | 7.3 | 9.9 | 10.6 | 6.9 | 7.0 | 2.7 | 2.6 |
| Person | 7.6 | 7.8 | 9.4 | 10.2 | 7.8 | 8.1 | 2.1 | 2.0 |
| Above 3 SD Overnourished (BMI for Age) (%) | | | | | | | | |
| Male | 4.0 | 4.3 | 4.4 | 4.4 | - | - | - | - |
| Female | 4.1 | 4.0 | 2.7 | 3.1 | - | - | - | - |
| Person | 4.0 | 4.1 | 3.6 | 3.8 | - | - | - | - |
| Nutritional status 5-18 years | | | | | | | | |
| Below -2 SD Undernourished (BMI for Age) (%) | | | | | | | | |
| Male | 33.3 | 34.7 | 29.9 | 30.9 | 26.5 | 26.8 | 45.7 | 47.0 |
| Female | 26.3 | 27.0 | 25.3 | 26.3 | 20.3 | 20.6 | 31.5 | 32.1 |
| Person | 29.9 | 30.9 | 27.7 | 28.6 | 23.6 | 23.9 | 38.5 | 39.4 |
| Below -3 SD Undernourished (BMI for Age) (%) | | | | | | | | |
| Male | 17.1 | 17.7 | 12.5 | 13.2 | 13.5 | 13.7 | 22.9 | 23.6 |
| Female | 12.6 | 12.8 | 7.7 | 7.0 | 7.3 | 7.3 | 12.0 | 12.1 |
| Person | 14.9 | 15.3 | 10.2 | 10.1 | 10.6 | 10.7 | 17.4 | 17.7 |
| Above 2 SD Overnourished (BMI for Age) (%) | | | | | | | | |
| Male | 1.7 | 1.6 | 1.9 | 1.5 | 1.4 | 1.5 | - | - |
| Female | 1.4 | 1.4 | 0.7 | 0.6 | 0.9 | - | - | - |
| Person | 1.5 | 1.5 | 1.3 | 1.0 | 1.1 | 1.2 | - | - |
| Above 3 SD Overnourished (BMI for Age) (%) | | | | | | | | |
| Male | 0.8 | 0.8 | 1.2 | - | - | - | - | - |
| Female | 0.6 | 0.6 | - | - | - | - | - | - |
| Person | 0.7 | 0.7 | 0.8 | - | 0.6 | 0.7 | - | - |

| Indicators | Chhattisgarh | | Koriya | | Surguja | | Jashpur | |
|--|--------------|-------|--------|-------|---------|-------|---------|-------|
| | Total | Rural | Total | Rural | Total | Rural | Total | Rural |
| Nutritional status 18 years and above | | | | | | | | |
| BMI ≥25.0 for age 18-59 year (%) | | | | | | | | |
| Male | 6.5 | 4.9 | 8.1 | 5.6 | 1.7 | 1.4 | 2.4 | 2.2 |
| Female | 6.2 | 5.0 | 11.7 | 7.0 | 1.0 | 0.8 | 2.7 | 2.7 |
| Person | 6.3 | 4.9 | 9.9 | 6.3 | 1.4 | 1.1 | 2.6 | 2.5 |
| BMI ≥25.0 for age 60 year and above (%) | | | | | | | | |
| Male | 7.4 | 5.2 | 9.0 | 9.1 | - | - | 3.5 | 3.4 |
| Female | 8.6 | 7.1 | 13.1 | 11.9 | - | - | 2.8 | 2.7 |
| Person | 8.0 | 6.1 | 10.9 | 10.4 | - | - | 3.2 | 3.1 |
| BMI ≥30.0 for age 18-59 year (%) | | | | | | | | |
| Male | 1.1 | 0.9 | 1.5 | 0.9 | 0.5 | - | - | - |
| Female | 1.4 | 1.2 | 2.6 | 1.2 | 0.2 | - | 0.5 | - |
| Person | 1.3 | 1.0 | 2.1 | 1.1 | 0.3 | 0.2 | 0.5 | - |
| BMI ≥30.0 for age 60 year and above (%) | | | | | | | | |
| Male | 0.9 | 0.8 | - | - | - | - | - | - |
| Female | 2.2 | 1.9 | 3.4 | 2.3 | - | - | - | - |
| Person | 1.5 | 1.3 | 2.2 | 1.8 | - | - | - | - |
| BMI <18.5 for age 18-59 year (%) | | | | | | | | |
| Male | 12.5 | 13.9 | 25.7 | 26.8 | 10.9 | 10.8 | 11.9 | 12.3 |
| Female | 18.5 | 20.7 | 26.9 | 31.7 | 21.2 | 21.6 | 19.1 | 19.4 |
| Person | 15.4 | 17.3 | 26.3 | 29.3 | 16.2 | 16.4 | 15.8 | 16.2 |
| BMI <18.5 for age 60 year and above (%) | | | | | | | | |
| Male | 22.9 | 25.8 | 36.2 | 37.4 | 29.7 | 30.8 | 29.0 | 30.0 |
| Female | 28.4 | 30.9 | 29.8 | 30.9 | 46.3 | 49.2 | 48.0 | 49.2 |
| Person | 25.6 | 28.3 | 33.3 | 34.5 | 37.9 | 40.0 | 38.0 | 39.1 |
| Anaemia status by Haemoglobin Level | | | | | | | | |
| Anaemia in the age group 6-59 months (%) | | | | | | | | |
| Male | 64.8 | 67.3 | 76.0 | 76.8 | 71.7 | 72.3 | 85.6 | 85.4 |
| Female | 62.7 | 65.9 | 68.7 | 68.7 | 64.6 | 65.5 | 84.2 | 84.0 |
| Person | 63.8 | 66.6 | 72.6 | 72.7 | 68.3 | 69.0 | 84.9 | 84.7 |
| Severe Anemia in the age group 6-59 months (%) | | | | | | | | |
| Male | 1.0 | 1.0 | - | - | - | - | 5.7 | 5.7 |
| Female | 0.9 | 1.1 | - | - | - | - | 7.4 | 7.2 |
| Person | 1.0 | 1.0 | - | - | - | - | 6.5 | 6.5 |
| Anaemia in the age group 5-9 years (%) | | | | | | | | |
| Male | 78.5 | 80.4 | 86.3 | 87.1 | 77.0 | 78.4 | 85.8 | 85.6 |
| Female | 78.4 | 80.0 | 88.6 | 90.1 | 77.6 | 78.2 | 85.9 | 85.6 |
| Person | 78.5 | 80.2 | 87.5 | 88.6 | 77.3 | 78.3 | 85.9 | 85.6 |
| Severe Anemia in the age group 5-9 years (%) | | | | | | | | |
| Male | 10.5 | 11.0 | 6.2 | - | 9.8 | 9.7 | 32.8 | 32.9 |
| Female | 10.9 | 11.0 | 9.5 | 11.6 | 7.6 | 7.4 | 35.1 | 35.5 |
| Person | 10.7 | 11.0 | 7.8 | 9.0 | 8.8 | 8.6 | 34.0 | 34.2 |
| Anaemia in the age group 10-17 years (%) | | | | | | | | |
| Male | 74.2 | 75.4 | 77.4 | 78.6 | 81.6 | 81.7 | 81.6 | 82.1 |
| Female | 76.0 | 77.3 | 81.1 | 81.6 | 81.4 | 82.0 | 88.3 | 88.4 |
| Person | 75.1 | 76.3 | 79.2 | 80.1 | 81.5 | 81.8 | 85.1 | 85.4 |
| Severe Anemia in the age group 10-17 years (%) | | | | | | | | |
| Male | 9.7 | 10.3 | 7.3 | 8.0 | 11.2 | 10.2 | 27.9 | 28.3 |
| Female | 9.3 | 9.8 | 10.4 | 11.3 | 8.2 | 8.1 | 30.6 | 30.1 |
| Person | 9.5 | 10.0 | 8.8 | 9.6 | 9.8 | 9.2 | 29.3 | 29.3 |
| Anaemia in the age group 18-59 years (%) | | | | | | | | |
| Male | 83.6 | 84.8 | 87.8 | 88.5 | 90.7 | 91.3 | 91.6 | 91.1 |
| Female | 76.4 | 77.9 | 79.9 | 82.0 | 81.1 | 82.5 | 87.1 | 86.8 |
| Severe Anemia in the age group 18-59 years (%) | | | | | | | | |
| Male | 8.8 | 9.3 | 6.3 | 6.6 | 11.1 | 10.6 | 23.1 | 22.0 |
| Female | 10.6 | 11.1 | 9.4 | 9.8 | 11.6 | 11.4 | 23.6 | 22.8 |
| Anaemia in the age group 60 years and above (%) | | | | | | | | |
| Male | 84.9 | 85.9 | 89.0 | 90.0 | 91.7 | 92.1 | 93.2 | 93.0 |
| Female | 78.1 | 78.5 | 84.0 | 82.6 | 77.7 | 78.0 | 90.6 | 90.5 |
| Severe Anemia in the age group 60 years and above (%) | | | | | | | | |
| Male | 9.9 | 10.5 | 6.8 | 7.6 | 8.7 | 8.1 | 18.7 | 17.7 |
| Female | 9.4 | 9.1 | 8.8 | 6.9 | 11.0 | 10.1 | 18.4 | 17.9 |

| Indicators | Chhattisgarh | | Koriya | | Surguja | | Jashpur | |
|--|--------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| | Total | Rural | Total | Rural | Total | Rural | Total | Rural |
| Blood Sugar Level (18 years and above) | | | | | | | | |
| Blood Sugar Level ≥110 mg/dl (%) | | | | | | | | |
| Male | 13.2 | 12.2 | 11.8 | 11.4 | 9.9 | 9.7 | 11.1 | 11.1 |
| Female | 9.9 | 9.3 | 11.5 | 9.3 | 6.4 | 6.2 | 9.1 | 9.1 |
| Person | 11.5 | 10.7 | 11.6 | 10.3 | 8.2 | 7.9 | 9.9 | 10.0 |
| Blood Sugar Level ≥130 mg/dl (%) | | | | | | | | |
| Male | 5.1 | 4.3 | 4.8 | 4.6 | 1.6 | 1.5 | 2.4 | 2.4 |
| Female | 3.6 | 3.2 | 4.6 | 3.4 | 1.0 | - | 2.3 | 2.2 |
| Person | 4.3 | 3.7 | 4.7 | 4.0 | 1.4 | 1.2 | 2.3 | 2.3 |
| Blood Sugar Level ≥150 mg/dl (%) | | | | | | | | |
| Male | 1.9 | 1.7 | 1.8 | 1.6 | 0.7 | 0.7 | 1.2 | 1.2 |
| Female | 1.5 | 1.3 | 2.3 | 1.4 | 0.6 | - | 0.9 | 0.8 |
| Person | 1.7 | 1.5 | 2.1 | 1.5 | 0.7 | 0.6 | 1.0 | 1.0 |
| Hypertension (18 years and above) | | | | | | | | |
| Above Normal Range | | | | | | | | |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%) | | | | | | | | |
| Male | 6.1 | 5.7 | 7.2 | 7.3 | 7.7 | 7.7 | 4.1 | 3.9 |
| Female | 6.1 | 6.0 | 7.5 | 6.5 | 4.6 | 4.5 | 3.3 | 3.3 |
| Person | 6.1 | 5.9 | 7.4 | 6.9 | 6.0 | 6.0 | 3.7 | 3.6 |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%) | | | | | | | | |
| Male | 5.3 | 4.9 | 7.3 | 7.5 | 5.3 | 5.0 | 3.0 | 2.8 |
| Female | 4.2 | 4.2 | 6.4 | 6.0 | 3.6 | 3.4 | 2.5 | 2.5 |
| Person | 4.8 | 4.5 | 6.8 | 6.8 | 4.4 | 4.1 | 2.7 | 2.7 |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%) | | | | | | | | |
| Male | 8.3 | 8.2 | 5.8 | 4.9 | 5.7 | 5.3 | 3.5 | 3.4 |
| Female | 6.5 | 6.4 | 5.0 | 3.5 | 4.4 | 4.3 | 2.6 | 2.5 |
| Person | 7.4 | 7.3 | 5.4 | 4.2 | 5.0 | 4.8 | 3.0 | 2.9 |
| Moderately high | | | | | | | | |
| Systolic ≥160mm of Hg & Diastolic ≥100mm of Hg (%) | | | | | | | | |
| Male | 1.8 | 1.7 | 2.6 | 2.5 | 1.1 | 1.0 | 0.7 | 0.7 |
| Female | 2.4 | 2.4 | 2.0 | 1.9 | 0.8 | 0.7 | 1.0 | 1.0 |
| Person | 2.1 | 2.0 | 2.3 | 2.2 | 0.9 | 0.9 | 0.9 | 0.9 |
| Systolic ≥160mm of Hg & Diastolic <100mm of Hg (%) | | | | | | | | |
| Male | 1.8 | 1.7 | 1.2 | 1.3 | 1.2 | 1.3 | 0.9 | 0.9 |
| Female | 1.9 | 1.8 | 2.4 | 2.1 | 1.1 | 1.0 | 0.9 | 1.0 |
| Person | 1.8 | 1.7 | 1.9 | 1.7 | 1.1 | 1.1 | 0.9 | 0.9 |
| Systolic <160mm of Hg & Diastolic ≥100mm of Hg (%) | | | | | | | | |
| Male | 3.1 | 2.9 | 2.8 | 2.6 | 1.6 | 1.6 | 1.1 | - |
| Female | 2.7 | 2.8 | 2.1 | 1.6 | 1.0 | 1.0 | 0.7 | 0.7 |
| Person | 2.9 | 2.8 | 2.5 | 2.1 | 1.3 | 1.3 | 0.9 | 0.8 |
| Very high | | | | | | | | |
| Systolic ≥180mm of Hg & Diastolic ≥110mm of Hg (%) | | | | | | | | |
| Male | 0.6 | 0.6 | 0.6 | 0.7 | 0.3 | 0.3 | - | - |
| Female | 0.8 | 0.8 | 0.7 | 0.8 | - | - | 0.4 | 0.4 |
| Person | 0.7 | 0.7 | 0.6 | 0.7 | 0.2 | 0.2 | 0.3 | 0.3 |
| Systolic ≥180mm of Hg & Diastolic <110mm of Hg (%) | | | | | | | | |
| Male | 0.5 | 0.5 | 0.8 | 0.9 | 0.4 | 0.4 | - | - |
| Female | 0.9 | 0.9 | 0.6 | 0.4 | 0.4 | 0.3 | 0.5 | 0.5 |
| Person | 0.7 | 0.7 | 0.7 | 0.6 | 0.4 | 0.4 | 0.3 | 0.3 |
| Systolic <180mm of Hg & Diastolic ≥110mm of Hg (%) | | | | | | | | |
| Male | 1.6 | 1.6 | 1.5 | 1.3 | 0.5 | 0.5 | 0.3 | - |
| Female | 1.6 | 1.6 | 0.8 | 0.7 | 0.3 | 0.2 | - | - |
| Person | 1.6 | 1.6 | 1.1 | 1.0 | 0.4 | 0.4 | 0.3 | 0.2 |
| Iodine content in Household Salt (more than 15 ppm) (%) | 75.9 | 75.6 | 70.5 | 74.7 | 64.5 | 62.6 | 64.7 | 65.3 |

| Indicators | Raigarh | | Korba | | Janjgir-Champa | | Bilaspur | |
|--|---------|-------|-------|-------|----------------|-------|----------|-------|
| | Total | Rural | Total | Rural | Total | Rural | Total | Rural |
| Nutritional status below 5 years | | | | | | | | |
| Below -2 SD Wasting (Weight for Height) (%) | | | | | | | | |
| Male | 23.2 | 23.0 | 23.5 | 28.7 | 28.8 | 31.0 | 33.2 | 30.2 |
| Female | 28.9 | 27.8 | 26.6 | 33.1 | 27.6 | 28.7 | 34.3 | 31.3 |
| Person | 26.0 | 25.2 | 24.8 | 30.5 | 28.3 | 30.0 | 33.7 | 30.7 |
| Below -3 SD Wasting (Weight for Height) (%) | | | | | | | | |
| Male | 7.3 | 6.8 | - | - | 5.9 | 6.0 | 14.5 | - |
| Female | 7.2 | 7.9 | - | - | - | 3.8 | 15.9 | 14.4 |
| Person | 7.3 | 7.3 | - | - | 4.4 | 5.0 | 15.1 | 14.9 |
| Below -2 SD Stunting (Height for Age) (%) | | | | | | | | |
| Male | 46.5 | 46.1 | 39.3 | 41.0 | 31.6 | 27.3 | 38.3 | 41.7 |
| Female | 40.0 | 42.0 | 25.4 | 28.4 | 31.6 | 29.3 | 34.2 | 39.6 |
| Person | 43.2 | 44.2 | 33.5 | 35.9 | 31.6 | 28.1 | 36.5 | 40.7 |
| Below -3SD Stunting (Height for Age) (%) | | | | | | | | |
| Male | 31.1 | 31.1 | 22.1 | 26.3 | 16.0 | 11.0 | 23.8 | 25.4 |
| Female | 29.8 | 32.5 | 13.4 | 18.4 | 20.5 | 16.5 | 24.5 | 30.8 |
| Person | 30.4 | 31.8 | 18.4 | 23.1 | 17.9 | 13.3 | 24.1 | 27.9 |
| Below -2 SD Underweight (Weight for Age) (%) | | | | | | | | |
| Male | 42.7 | 42.8 | 39.2 | 42.8 | 30.7 | 25.8 | 45.5 | 43.5 |
| Female | 33.6 | 35.1 | 27.5 | 35.8 | 34.5 | 27.0 | 53.6 | 53.7 |
| Person | 38.4 | 39.2 | 34.3 | 39.9 | 32.4 | 26.3 | 49.1 | 48.3 |
| Below -3 SD Underweight (Weight for Age) (%) | | | | | | | | |
| Male | 23.2 | 23.6 | 13.6 | 12.0 | 18.4 | 11.8 | 31.1 | 28.2 |
| Female | 14.8 | 16.1 | 7.8 | 14.6 | 20.4 | 11.5 | 28.4 | 27.5 |
| Person | 19.2 | 20.1 | 11.2 | 13.1 | 19.3 | 11.7 | 29.9 | 27.9 |
| Below -2 SD Undernourished (BMI for Age) (%) | | | | | | | | |
| Male | 19.4 | 19.1 | 20.8 | 24.6 | 31.4 | 34.0 | 34.6 | 27.7 |
| Female | 27.9 | 26.0 | 24.2 | 29.6 | 34.5 | 32.6 | 35.2 | 28.5 |
| Person | 23.3 | 22.2 | 22.3 | 26.7 | 32.8 | 33.4 | 34.9 | 28.1 |
| Below -3 SD Undernourished (BMI for Age) (%) | | | | | | | | |
| Male | 6.7 | 6.2 | 4.6 | - | 11.2 | 11.8 | 23.3 | 17.7 |
| Female | 11.9 | 11.5 | - | - | 8.4 | - | 24.1 | 17.8 |
| Person | 9.1 | 8.6 | 3.3 | - | 10.0 | 10.0 | 23.7 | 17.7 |
| Above 2 SD Overnourished (BMI for Age) (%) | | | | | | | | |
| Male | 17.3 | 17.0 | 3.9 | 5.2 | 7.5 | 6.0 | 11.4 | 13.3 |
| Female | 10.7 | 11.6 | 5.5 | - | 6.4 | 6.8 | 10.7 | 13.5 |
| Person | 14.2 | 14.5 | 4.6 | 5.4 | 7.0 | 6.4 | 11.1 | 13.4 |
| Above 3 SD Overnourished (BMI for Age) (%) | | | | | | | | |
| Male | 7.1 | 7.2 | 1.6 | - | 3.3 | - | 5.4 | 6.2 |
| Female | - | - | 4.1 | - | - | - | - | - |
| Person | 6.7 | 7.0 | 2.7 | - | 3.6 | 3.0 | 5.1 | 6.1 |
| Nutritional status 5-18 years | | | | | | | | |
| Below -2 SD Undernourished (BMI for Age) (%) | | | | | | | | |
| Male | 23.1 | 22.6 | 32.5 | 34.9 | 29.8 | 31.0 | 26.1 | 29.4 |
| Female | 17.3 | 16.8 | 28.7 | 35.0 | 20.9 | 20.4 | 24.3 | 28.1 |
| Person | 20.1 | 19.7 | 30.6 | 34.9 | 25.3 | 25.8 | 25.3 | 28.8 |
| Below -3 SD Undernourished (BMI for Age) (%) | | | | | | | | |
| Male | 9.7 | 8.8 | 13.9 | 14.0 | 12.6 | 12.2 | 14.7 | 16.3 |
| Female | 8.4 | 8.4 | 10.5 | 10.8 | 6.3 | 5.5 | 13.5 | 17.6 |
| Person | 9.0 | 8.6 | 12.2 | 12.4 | 9.4 | 8.9 | 14.2 | 16.9 |
| Above 2 SD Overnourished (BMI for Age) (%) | | | | | | | | |
| Male | 1.1 | 1.3 | 2.1 | 1.7 | 1.0 | - | 3.7 | 4.8 |
| Female | - | - | 1.2 | - | - | - | - | - |
| Person | 1.0 | 1.1 | 1.7 | 1.5 | 0.7 | 0.7 | 3.5 | 4.6 |
| Above 3 SD Overnourished (BMI for Age) (%) | | | | | | | | |
| Male | - | - | 1.0 | - | 0.5 | - | - | - |
| Female | - | - | - | - | - | - | - | - |
| Person | 0.4 | 0.4 | 0.7 | 0.3 | 0.3 | 0.3 | - | - |

| Indicators | Raigarh | | Korba | | Janjgir-Champa | | Bilaspur | |
|--|---------|-------|-------|-------|----------------|-------|----------|-------|
| | Total | Rural | Total | Rural | Total | Rural | Total | Rural |
| Nutritional status 18 years and above | | | | | | | | |
| BMI ≥25.0 for age 18-59 year (%) | | | | | | | | |
| Male | 5.8 | 2.4 | 14.0 | 6.7 | 7.6 | 6.2 | 5.5 | 6.2 |
| Female | 4.9 | 3.4 | 11.8 | 4.5 | 5.7 | 5.0 | 5.5 | 5.9 |
| Person | 5.3 | 2.9 | 12.9 | 5.6 | 6.7 | 5.6 | 5.5 | 6.0 |
| BMI ≥25.0 for age 60 year and above (%) | | | | | | | | |
| Male | 14.3 | 4.8 | - | - | 4.9 | 3.0 | 5.5 | 5.3 |
| Female | 7.3 | - | 15.2 | 6.6 | 5.6 | 4.0 | 5.6 | 5.5 |
| Person | 10.6 | 3.9 | 11.7 | 4.2 | 5.2 | 3.5 | 5.5 | 5.4 |
| BMI ≥30.0 for age 18-59 year (%) | | | | | | | | |
| Male | - | - | 2.4 | - | 1.0 | - | 0.9 | 1.2 |
| Female | 0.9 | 0.7 | 3.0 | - | 0.7 | 0.5 | 1.2 | 1.4 |
| Person | 0.6 | 0.4 | 2.7 | - | 0.8 | 0.6 | 1.0 | 1.3 |
| BMI ≥30.0 for age 60 year and above (%) | | | | | | | | |
| Male | - | - | - | - | - | - | - | - |
| Female | - | - | - | - | 1.6 | - | 0.4 | 0.5 |
| Person | - | - | - | - | 1.2 | 0.9 | 0.5 | 0.7 |
| BMI <18.5 for age 18-59 year (%) | | | | | | | | |
| Male | 12.3 | 13.6 | 12.2 | 14.6 | 11.8 | 12.5 | 7.0 | 9.7 |
| Female | 12.4 | 12.9 | 17.6 | 22.4 | 17.8 | 19.0 | 11.8 | 15.2 |
| Person | 12.3 | 13.2 | 14.9 | 18.4 | 14.7 | 15.6 | 9.3 | 12.3 |
| BMI <18.5 for age 60 year and above (%) | | | | | | | | |
| Male | 23.3 | 27.7 | 24.9 | 40.3 | 25.5 | 26.8 | 22.2 | 27.5 |
| Female | 22.5 | 25.2 | 25.7 | 30.3 | 32.9 | 34.4 | 31.3 | 38.6 |
| Person | 22.8 | 26.3 | 25.3 | 35.6 | 29.3 | 30.7 | 26.6 | 33.1 |
| Anaemia status by Haemoglobin Level | | | | | | | | |
| Anaemia in the age group 6-59 months (%) | | | | | | | | |
| Male | 70.9 | 70.1 | 67.1 | 76.8 | 63.0 | 59.7 | 63.3 | 61.4 |
| Female | 78.8 | 75.9 | 63.5 | 62.6 | 62.4 | 63.0 | 64.3 | 66.5 |
| Person | 75.1 | 73.1 | 65.5 | 70.4 | 62.7 | 61.1 | 63.8 | 63.9 |
| Severe Anemia in the age group 6-59 months (%) | | | | | | | | |
| Male | - | - | - | - | - | - | - | - |
| Female | - | - | - | - | - | - | - | - |
| Person | - | - | - | - | - | - | - | - |
| Anaemia in the age group 5-9 years (%) | | | | | | | | |
| Male | 87.0 | 86.4 | 76.7 | 86.1 | 75.8 | 75.8 | 77.2 | 79.5 |
| Female | 86.3 | 86.2 | 84.0 | 84.9 | 72.9 | 71.0 | 71.0 | 75.6 |
| Person | 86.6 | 86.3 | 80.4 | 85.5 | 74.5 | 73.6 | 74.4 | 77.7 |
| Severe Anemia in the age group 5-9 years (%) | | | | | | | | |
| Male | 14.2 | 14.3 | 9.2 | 9.9 | - | - | 4.0 | - |
| Female | 11.9 | 11.3 | 7.5 | 10.5 | 4.9 | 4.7 | 6.6 | 6.1 |
| Person | 13.1 | 12.8 | 8.3 | 10.2 | 3.2 | 3.2 | 5.1 | 4.2 |
| Anaemia in the age group 10-17 years (%) | | | | | | | | |
| Male | 84.6 | 84.0 | 77.4 | 80.1 | 65.2 | 66.0 | 69.3 | 72.5 |
| Female | 78.0 | 77.1 | 79.9 | 85.5 | 73.7 | 72.8 | 64.8 | 66.4 |
| Person | 81.4 | 80.5 | 78.6 | 82.8 | 69.7 | 69.5 | 67.2 | 69.7 |
| Severe Anemia in the age group 10-17 years (%) | | | | | | | | |
| Male | 13.1 | 14.0 | 5.8 | - | 2.2 | 2.1 | 5.0 | 3.8 |
| Female | 9.2 | 9.4 | 8.2 | 9.7 | 5.8 | 5.4 | 3.0 | 2.8 |
| Person | 11.2 | 11.7 | 7.0 | 8.2 | 4.1 | 3.8 | 4.1 | 3.4 |
| Anaemia in the age group 18-59 years (%) | | | | | | | | |
| Male | 90.5 | 90.3 | 78.5 | 82.8 | 74.1 | 74.6 | 77.6 | 77.0 |
| Female | 81.2 | 80.9 | 76.3 | 80.2 | 74.2 | 72.5 | 60.9 | 60.5 |
| Severe Anemia in the age group 18-59 years (%) | | | | | | | | |
| Male | 10.7 | 10.2 | 5.2 | 7.4 | 3.9 | 4.3 | 3.9 | 3.3 |
| Female | 9.7 | 10.1 | 7.0 | 7.2 | 5.2 | 5.3 | 4.5 | 2.8 |
| Anaemia in the age group 60 years and above (%) | | | | | | | | |
| Male | 93.5 | 95.0 | 83.5 | 88.8 | 79.7 | 79.5 | 73.1 | 70.9 |
| Female | 81.0 | 82.2 | 74.9 | 76.5 | 74.7 | 72.3 | 66.0 | 67.0 |
| Severe Anemia in the age group 60 years and above (%) | | | | | | | | |
| Male | 7.3 | 7.2 | 13.0 | 16.0 | 2.6 | 3.0 | 4.0 | - |
| Female | 10.2 | 10.3 | 7.5 | 9.1 | 5.9 | 5.2 | 6.0 | - |

| Indicators | Raigarh | | Korba | | Janjgir-Champa | | Bilaspur | |
|--|-------------|-------------|-------------|-------------|----------------|-------------|-------------|-------------|
| | Total | Rural | Total | Rural | Total | Rural | Total | Rural |
| Blood Sugar Level (18 years and above) | | | | | | | | |
| Blood Sugar Level ≥110 mg/dl (%) | | | | | | | | |
| Male | 16.8 | 15.4 | 10.4 | 2.8 | 5.8 | 5.2 | 9.3 | 9.1 |
| Female | 11.1 | 10.3 | 6.1 | 4.1 | 5.2 | 4.8 | 8.5 | 10.2 |
| Person | 14.1 | 12.8 | 8.4 | 3.4 | 5.5 | 5.0 | 8.9 | 9.6 |
| Blood Sugar Level ≥130 mg/dl (%) | | | | | | | | |
| Male | 7.4 | 7.6 | 4.1 | 1.4 | 2.9 | 2.2 | 3.5 | 4.0 |
| Female | 3.4 | 3.5 | 3.1 | 3.0 | 2.5 | 2.2 | 2.7 | 3.7 |
| Person | 5.5 | 5.5 | 3.7 | 2.2 | 2.7 | 2.2 | 3.1 | 3.8 |
| Blood Sugar Level ≥150 mg/dl (%) | | | | | | | | |
| Male | 2.5 | 2.6 | 2.1 | 0.8 | 2.3 | 1.7 | 2.8 | - |
| Female | 1.0 | 1.1 | 2.0 | - | 1.8 | 1.5 | 2.1 | 2.9 |
| Person | 1.8 | 1.8 | 2.1 | 1.4 | 2.0 | 1.6 | 2.5 | 3.1 |
| Hypertension (18 years and above) | | | | | | | | |
| Above Normal Range | | | | | | | | |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%) | | | | | | | | |
| Male | 4.2 | 4.3 | 4.7 | 3.1 | 6.1 | 5.9 | 4.8 | 6.6 |
| Female | 3.2 | 3.6 | 4.7 | 3.2 | 6.2 | 5.3 | 6.2 | 7.9 |
| Person | 3.7 | 3.9 | 4.7 | 3.1 | 6.1 | 5.6 | 5.5 | 7.2 |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%) | | | | | | | | |
| Male | 5.7 | 5.7 | 3.0 | 2.9 | 5.2 | 4.4 | 4.9 | 4.7 |
| Female | 4.0 | 4.3 | 2.5 | - | 4.8 | 4.3 | 3.4 | 3.9 |
| Person | 4.9 | 5.0 | 2.7 | 2.8 | 5.0 | 4.4 | 4.2 | 4.3 |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%) | | | | | | | | |
| Male | 8.1 | 8.0 | 8.9 | 10.4 | 9.2 | 9.6 | 7.0 | 9.3 |
| Female | 4.9 | 4.6 | 6.6 | 7.8 | 7.7 | 7.9 | 6.9 | 9.3 |
| Person | 6.5 | 6.3 | 7.7 | 9.2 | 8.5 | 8.7 | 7.0 | 9.3 |
| Moderately high | | | | | | | | |
| Systolic ≥160mm of Hg & Diastolic ≥100mm of Hg (%) | | | | | | | | |
| Male | 0.5 | 0.5 | 1.0 | - | 3.0 | 2.6 | 3.0 | 4.1 |
| Female | 0.7 | 0.7 | 1.2 | 0.7 | 2.6 | 2.1 | 4.2 | 5.4 |
| Person | 0.6 | 0.6 | 1.1 | 0.6 | 2.8 | 2.3 | 3.6 | 4.7 |
| Systolic ≥160mm of Hg & Diastolic <100mm of Hg (%) | | | | | | | | |
| Male | 0.7 | 0.8 | - | - | 2.2 | 2.2 | 2.7 | 3.4 |
| Female | 0.8 | 0.9 | 1.7 | 1.2 | 2.8 | 2.4 | 1.9 | 2.3 |
| Person | 0.8 | 0.9 | 1.1 | 0.7 | 2.5 | 2.3 | 2.3 | 2.8 |
| Systolic <160mm of Hg & Diastolic ≥100mm of Hg (%) | | | | | | | | |
| Male | 3.5 | 3.7 | 1.6 | 0.9 | 2.6 | 2.5 | 7.1 | 9.9 |
| Female | 1.9 | 1.9 | 1.8 | 1.5 | 2.2 | 2.1 | 7.9 | 10.5 |
| Person | 2.7 | 2.8 | 1.7 | 1.2 | 2.4 | 2.3 | 7.5 | 10.2 |
| Very high | | | | | | | | |
| Systolic ≥180mm of Hg & Diastolic ≥110mm of Hg (%) | | | | | | | | |
| Male | - | - | - | - | 1.0 | 0.9 | 1.2 | 1.8 |
| Female | - | - | 0.6 | 0.4 | 0.9 | 0.7 | 2.0 | 2.7 |
| Person | - | - | 0.4 | 0.2 | 1.0 | 0.8 | 1.6 | 2.3 |
| Systolic ≥180mm of Hg & Diastolic <110mm of Hg (%) | | | | | | | | |
| Male | - | - | - | - | 0.9 | 1.1 | 0.6 | - |
| Female | 0.6 | 0.6 | - | - | 1.3 | 1.2 | 1.0 | 1.4 |
| Person | 0.3 | 0.4 | - | - | 1.1 | 1.1 | 0.8 | 1.0 |
| Systolic <180mm of Hg & Diastolic ≥110mm of Hg (%) | | | | | | | | |
| Male | - | - | 0.7 | - | 1.6 | 1.7 | 5.3 | 7.5 |
| Female | 0.7 | - | 0.6 | - | 0.8 | 0.8 | 6.1 | 8.3 |
| Person | 0.5 | 0.5 | 0.7 | 0.5 | 1.2 | 1.2 | 5.7 | 7.9 |
| Iodine content in Household Salt (more than 15 ppm) (%) | | | | | | | | |
| | 89.7 | 88.7 | 69.3 | 72.2 | 69.0 | 65.2 | 80.5 | 78.8 |

| Indicators | Kawardha | | Rajnandgaon | | Durg | | Raipur | |
|--|----------|-------|-------------|-------|-------|-------|--------|-------|
| | Total | Rural | Total | Rural | Total | Rural | Total | Rural |
| Nutritional status below 5 years | | | | | | | | |
| Below -2 SD Wasting (Weight for Height) (%) | | | | | | | | |
| Male | 31.3 | 25.9 | 24.0 | 25.8 | 36.3 | 34.4 | 29.7 | 22.5 |
| Female | 31.4 | 26.5 | 41.0 | 47.2 | 31.3 | 30.9 | 34.8 | 30.5 |
| Person | 31.3 | 26.2 | 32.5 | 36.5 | 34.1 | 32.9 | 32.5 | 26.9 |
| Below -3 SD Wasting (Weight for Height) (%) | | | | | | | | |
| Male | 18.8 | 14.1 | 7.3 | 8.8 | 5.6 | 4.4 | 13.6 | 5.1 |
| Female | 10.3 | - | - | - | - | - | 12.0 | 6.2 |
| Person | 15.2 | 10.8 | 4.5 | 5.4 | 6.5 | 6.5 | 12.8 | 5.7 |
| Below -2 SD Stunting (Height for Age) (%) | | | | | | | | |
| Male | 50.5 | 53.0 | 61.7 | 65.2 | 29.0 | 29.0 | 53.8 | 65.0 |
| Female | 33.1 | 34.7 | 46.6 | 46.3 | 36.0 | 38.0 | 48.9 | 55.2 |
| Person | 43.0 | 44.9 | 54.0 | 55.5 | 32.1 | 32.6 | 51.1 | 59.6 |
| Below -3SD Stunting (Height for Age) (%) | | | | | | | | |
| Male | 30.2 | 32.5 | 38.5 | 39.8 | 14.4 | 14.7 | 40.3 | 46.4 |
| Female | 20.7 | 22.7 | 27.6 | 29.0 | 17.1 | 11.9 | 29.4 | 27.8 |
| Person | 26.1 | 28.2 | 33.0 | 34.3 | 15.6 | 13.5 | 34.3 | 36.2 |
| Below -2 SD Underweight (Weight for Age) (%) | | | | | | | | |
| Male | 58.9 | 54.2 | 48.1 | 52.1 | 30.7 | 30.9 | 57.4 | 68.4 |
| Female | 40.4 | 36.4 | 44.9 | 48.0 | 32.3 | 28.5 | 47.8 | 56.7 |
| Person | 51.1 | 46.6 | 46.6 | 50.1 | 31.5 | 30.0 | 52.3 | 62.0 |
| Below -3 SD Underweight (Weight for Age) (%) | | | | | | | | |
| Male | 33.3 | 29.8 | 26.2 | 27.3 | 10.2 | 9.3 | 26.7 | 27.7 |
| Female | 13.4 | - | 25.9 | 27.7 | 11.7 | 8.7 | 26.9 | 31.7 |
| Person | 24.9 | 20.7 | 26.1 | 27.5 | 10.9 | 9.1 | 26.8 | 29.9 |
| Below -2 SD Undernourished (BMI for Age) (%) | | | | | | | | |
| Male | 33.2 | 26.3 | 18.5 | 19.6 | 37.4 | 35.8 | 30.1 | 20.6 |
| Female | 31.4 | 24.7 | 36.8 | 42.9 | 31.7 | 31.7 | 34.2 | 28.8 |
| Person | 32.5 | 25.6 | 27.4 | 30.7 | 34.8 | 34.1 | 32.3 | 25.1 |
| Below -3 SD Undernourished (BMI for Age) (%) | | | | | | | | |
| Male | 23.9 | 16.3 | - | - | 11.3 | 11.4 | 19.3 | 7.5 |
| Female | 15.7 | 7.9 | - | - | 12.6 | 15.2 | 15.8 | 8.2 |
| Person | 20.5 | 12.7 | 2.9 | 3.4 | 11.9 | 13.0 | 17.4 | 7.9 |
| Above 2 SD Overnourished (BMI for Age) (%) | | | | | | | | |
| Male | 12.0 | 11.6 | 23.0 | 25.4 | 3.6 | 4.0 | 5.5 | - |
| Female | 8.7 | 10.4 | 14.6 | 13.9 | 6.3 | 3.9 | 6.7 | - |
| Person | 10.6 | 11.1 | 19.0 | 19.9 | 4.8 | 4.0 | 6.1 | 5.1 |
| Above 3 SD Overnourished (BMI for Age) (%) | | | | | | | | |
| Male | 4.4 | 3.6 | 18.4 | 20.7 | 2.5 | - | 2.0 | - |
| Female | 4.8 | 5.6 | 11.9 | 11.4 | 4.5 | - | 3.0 | - |
| Person | 4.6 | 4.4 | 15.3 | 16.2 | 3.4 | 2.2 | 2.5 | 3.0 |
| Nutritional status 5-18 years | | | | | | | | |
| Below -2 SD Undernourished (BMI for Age) (%) | | | | | | | | |
| Male | 32.9 | 32.9 | 34.2 | 35.4 | 44.5 | 47.3 | 35.6 | 37.5 |
| Female | 26.8 | 27.2 | 28.4 | 28.4 | 33.4 | 35.0 | 28.5 | 27.7 |
| Person | 29.8 | 30.1 | 31.2 | 31.8 | 38.7 | 40.8 | 32.3 | 32.9 |
| Below -3 SD Undernourished (BMI for Age) (%) | | | | | | | | |
| Male | 20.3 | 21.6 | 16.7 | 17.2 | 22.5 | 23.4 | 22.5 | 24.4 |
| Female | 15.1 | 15.8 | 15.3 | 16.1 | 17.2 | 17.0 | 17.6 | 18.0 |
| Person | 17.7 | 18.7 | 16.0 | 16.6 | 19.8 | 20.0 | 20.2 | 21.4 |
| Above 2 SD Overnourished (BMI for Age) (%) | | | | | | | | |
| Male | - | - | 4.4 | - | 2.1 | 1.3 | 1.0 | - |
| Female | - | - | 4.6 | 5.1 | 1.4 | - | 0.9 | - |
| Person | - | - | 4.5 | 4.7 | 1.7 | 1.4 | 0.9 | 0.9 |
| Above 3 SD Overnourished (BMI for Age) (%) | | | | | | | | |
| Male | - | - | 1.5 | - | - | - | - | - |
| Female | - | - | 1.3 | 1.3 | - | - | 0.6 | - |
| Person | - | - | 1.4 | 1.4 | - | - | 0.5 | 0.5 |

| Indicators | Kawardha | | Rajnandgaon | | Durg | | Raipur | |
|--|----------|-------|-------------|-------|-------|-------|--------|-------|
| | Total | Rural | Total | Rural | Total | Rural | Total | Rural |
| Nutritional status 18 years and above | | | | | | | | |
| BMI ≥25.0 for age 18-59 year (%) | | | | | | | | |
| Male | 3.8 | 3.3 | 5.0 | 3.8 | 10.1 | 6.2 | 7.9 | 7.2 |
| Female | 2.3 | 1.9 | 6.7 | 5.4 | 9.3 | 8.3 | 7.7 | 6.9 |
| Person | 3.1 | 2.6 | 5.9 | 4.6 | 9.7 | 7.3 | 7.8 | 7.1 |
| BMI ≥25.0 for age 60 year and above (%) | | | | | | | | |
| Male | 5.1 | 5.0 | 7.5 | 5.2 | 14.1 | 8.2 | 8.3 | 7.2 |
| Female | 4.8 | 5.3 | 15.3 | 12.1 | 14.4 | 12.3 | 10.3 | 9.6 |
| Person | 4.9 | 5.2 | 11.6 | 8.8 | 14.3 | 10.3 | 9.2 | 8.4 |
| BMI ≥30.0 for age 18-59 year (%) | | | | | | | | |
| Male | - | - | - | - | 2.3 | 2.1 | 1.6 | - |
| Female | 0.6 | 0.7 | 1.8 | 1.8 | 2.5 | 2.7 | 2.2 | 2.1 |
| Person | 0.5 | 0.6 | 1.3 | 1.3 | 2.4 | 2.4 | 1.9 | 1.7 |
| BMI ≥30.0 for age 60 year and above (%) | | | | | | | | |
| Male | - | - | - | - | - | - | - | - |
| Female | - | - | 4.5 | - | 3.8 | - | 3.6 | - |
| Person | - | - | 2.5 | - | 2.7 | - | 2.3 | 2.5 |
| BMI <18.5 for age 18-59 year (%) | | | | | | | | |
| Male | 4.5 | 4.6 | 19.1 | 20.5 | 17.9 | 22.6 | 6.4 | 7.1 |
| Female | 6.5 | 7.2 | 24.1 | 26.5 | 23.3 | 29.3 | 10.8 | 11.7 |
| Person | 5.4 | 5.9 | 21.6 | 23.4 | 20.7 | 26.1 | 8.4 | 9.2 |
| BMI <18.5 for age 60 year and above (%) | | | | | | | | |
| Male | 9.5 | 10.6 | 21.6 | 24.6 | 21.5 | 26.6 | 13.1 | 15.1 |
| Female | 13.8 | 14.5 | 20.6 | 22.9 | 29.7 | 33.8 | 15.5 | - |
| Person | 11.6 | 12.5 | 21.1 | 23.7 | 25.7 | 30.4 | 14.2 | 15.2 |
| Anaemia status by Haemoglobin Level | | | | | | | | |
| Anaemia in the age group 6-59 months (%) | | | | | | | | |
| Male | 56.1 | 56.6 | 58.3 | 55.0 | 58.9 | 71.3 | 55.6 | 52.2 |
| Female | 55.3 | 53.2 | 66.9 | 62.5 | 54.9 | 73.3 | 54.6 | 56.2 |
| Person | 55.7 | 55.0 | 62.6 | 58.6 | 57.0 | 72.2 | 55.1 | 54.4 |
| Severe Anemia in the age group 6-59 months (%) | | | | | | | | |
| Male | - | - | - | - | - | - | - | - |
| Female | - | - | - | - | - | - | - | - |
| Person | - | - | - | - | - | - | - | - |
| Anaemia in the age group 5-9 years (%) | | | | | | | | |
| Male | 75.4 | 74.7 | 80.5 | 80.3 | 69.2 | 73.2 | 71.1 | 74.6 |
| Female | 73.5 | 72.8 | 79.4 | 78.0 | 67.8 | 76.6 | 75.0 | 74.0 |
| Person | 74.3 | 73.6 | 79.9 | 79.0 | 68.4 | 75.0 | 73.0 | 74.3 |
| Severe Anemia in the age group 5-9 years (%) | | | | | | | | |
| Male | 4.1 | 4.0 | 10.7 | 10.2 | 8.2 | 10.1 | 8.1 | 7.4 |
| Female | - | - | 8.4 | 6.6 | 6.9 | 8.7 | 11.0 | 9.1 |
| Person | 4.5 | 4.7 | 9.5 | 8.3 | 7.5 | 9.4 | 9.5 | 8.2 |
| Anaemia in the age group 10-17 years (%) | | | | | | | | |
| Male | 70.2 | 68.3 | 65.3 | 62.9 | 63.6 | 73.5 | 69.7 | 66.4 |
| Female | 80.2 | 80.1 | 75.3 | 73.4 | 65.7 | 73.1 | 71.7 | 70.7 |
| Person | 74.9 | 73.8 | 70.3 | 68.1 | 64.7 | 73.3 | 70.6 | 68.4 |
| Severe Anemia in the age group 10-17 years (%) | | | | | | | | |
| Male | 6.2 | 6.7 | 7.7 | 6.7 | 6.2 | 8.0 | 6.7 | 7.5 |
| Female | 7.5 | 8.0 | 8.4 | 6.0 | 6.9 | 7.8 | 6.7 | 7.7 |
| Person | 6.8 | 7.3 | 8.0 | 6.3 | 6.6 | 7.9 | 6.7 | 7.6 |
| Anaemia in the age group 18-59 years (%) | | | | | | | | |
| Male | 86.1 | 85.7 | 76.2 | 73.4 | 73.0 | 81.2 | 87.0 | 86.4 |
| Female | 79.9 | 79.8 | 77.9 | 75.2 | 69.9 | 75.5 | 73.3 | 74.3 |
| Severe Anemia in the age group 18-59 years (%) | | | | | | | | |
| Male | 6.7 | 6.8 | 6.9 | 6.1 | 4.8 | 5.9 | 7.6 | 7.7 |
| Female | 8.0 | 8.3 | 12.0 | 10.5 | 6.9 | 8.3 | 9.2 | 9.5 |
| Anaemia in the age group 60 years and above (%) | | | | | | | | |
| Male | 91.2 | 91.6 | 82.4 | 81.0 | 78.4 | 84.0 | 84.4 | 84.1 |
| Female | 77.8 | 78.1 | 83.9 | 82.4 | 78.1 | 81.0 | 74.0 | 72.8 |
| Severe Anemia in the age group 60 years and above (%) | | | | | | | | |
| Male | 12.5 | 13.7 | 16.4 | 17.1 | 10.2 | 11.7 | 5.0 | - |
| Female | 9.9 | 9.6 | 9.7 | - | 6.9 | 6.2 | 6.8 | 7.6 |

| Indicators | Kawardha | | Rajnandgaon | | Durg | | Raipur | |
|--|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| | Total | Rural | Total | Rural | Total | Rural | Total | Rural |
| Blood Sugar Level (18 years and above) | | | | | | | | |
| Blood Sugar Level ≥110 mg/dl (%) | | | | | | | | |
| Male | 18.1 | 17.7 | 19.7 | 20.5 | 15.4 | 12.7 | 11.7 | 7.9 |
| Female | 12.3 | 12.1 | 16.9 | 17.0 | 10.5 | 9.5 | 8.5 | 5.8 |
| Person | 15.3 | 14.9 | 18.2 | 18.7 | 12.8 | 11.1 | 10.2 | 6.9 |
| Blood Sugar Level ≥130 mg/dl (%) | | | | | | | | |
| Male | 6.0 | 6.3 | 8.2 | 8.3 | 9.6 | 7.0 | 5.4 | 3.2 |
| Female | 4.8 | 4.9 | 7.2 | 6.7 | 4.5 | 4.0 | 4.7 | 3.2 |
| Person | 5.4 | 5.6 | 7.7 | 7.5 | 6.9 | 5.5 | 5.1 | 3.2 |
| Blood Sugar Level ≥150 mg/dl (%) | | | | | | | | |
| Male | 0.5 | 0.6 | 2.7 | 2.4 | 3.6 | 3.6 | 1.4 | 0.6 |
| Female | - | - | 2.5 | 2.2 | 1.8 | 1.4 | 1.8 | 1.2 |
| Person | 0.5 | 0.5 | 2.6 | 2.3 | 2.6 | 2.5 | 1.6 | 0.9 |
| Hypertension (18 years and above) | | | | | | | | |
| Above Normal Range | | | | | | | | |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%) | | | | | | | | |
| Male | 9.0 | 9.9 | 11.2 | 10.7 | 9.6 | 7.8 | 4.4 | 2.8 |
| Female | 6.0 | 5.8 | 10.0 | 9.4 | 11.1 | 11.8 | 4.4 | 3.7 |
| Person | 7.5 | 7.9 | 10.5 | 10.0 | 10.4 | 9.9 | 4.4 | 3.2 |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%) | | | | | | | | |
| Male | 5.8 | 5.5 | 4.2 | 3.9 | 8.2 | 7.1 | 4.6 | 3.9 |
| Female | 2.5 | 2.3 | 7.2 | 7.2 | 5.8 | 6.1 | 3.3 | 2.8 |
| Person | 4.2 | 3.9 | 5.8 | 5.6 | 6.9 | 6.6 | 4.0 | 3.4 |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%) | | | | | | | | |
| Male | 8.7 | 8.4 | 9.7 | 9.9 | 7.3 | 6.4 | 9.7 | 8.5 |
| Female | 6.5 | 5.9 | 8.9 | 9.3 | 5.7 | 5.3 | 5.9 | 5.1 |
| Person | 7.6 | 7.2 | 9.3 | 9.6 | 6.5 | 5.8 | 7.9 | 6.8 |
| Moderately high | | | | | | | | |
| Systolic ≥160mm of Hg & Diastolic ≥100mm of Hg (%) | | | | | | | | |
| Male | 2.0 | 2.4 | 1.8 | 1.9 | 3.7 | 3.6 | 1.5 | 1.1 |
| Female | 1.7 | 1.7 | 2.5 | 2.5 | 5.9 | 6.5 | 2.4 | 2.1 |
| Person | 1.9 | 2.0 | 2.2 | 2.2 | 4.9 | 5.1 | 1.9 | 1.6 |
| Systolic ≥160mm of Hg & Diastolic <100mm of Hg (%) | | | | | | | | |
| Male | - | - | 2.6 | 2.5 | 2.7 | 2.9 | 2.6 | 1.8 |
| Female | 1.0 | 1.0 | 3.4 | 3.2 | 2.9 | 3.3 | 2.5 | 1.9 |
| Person | 0.9 | 0.9 | 3.0 | 2.9 | 2.8 | 3.1 | 2.5 | 1.8 |
| Systolic <160mm of Hg & Diastolic ≥100mm of Hg (%) | | | | | | | | |
| Male | 3.0 | 3.3 | 3.5 | 3.3 | 4.5 | 2.9 | 2.0 | - |
| Female | 2.1 | 2.0 | 3.2 | 3.3 | 3.9 | 3.8 | 2.0 | 2.1 |
| Person | 2.5 | 2.7 | 3.3 | 3.3 | 4.2 | 3.4 | 2.0 | 1.5 |
| Very high | | | | | | | | |
| Systolic ≥180mm of Hg & Diastolic ≥110mm of Hg (%) | | | | | | | | |
| Male | - | - | 0.7 | 0.8 | 1.3 | 1.3 | 0.6 | 0.5 |
| Female | - | - | 0.9 | 0.9 | 1.8 | 1.9 | 0.7 | - |
| Person | - | - | 0.8 | 0.8 | 1.6 | 1.6 | 0.6 | 0.5 |
| Systolic ≥180mm of Hg & Diastolic <110mm of Hg (%) | | | | | | | | |
| Male | - | - | 0.8 | - | 1.2 | 1.3 | 0.4 | 0.3 |
| Female | - | - | 1.3 | - | 1.8 | 2.1 | 1.0 | 0.8 |
| Person | - | - | 1.1 | 1.0 | 1.5 | 1.8 | 0.7 | 0.5 |
| Systolic <180mm of Hg & Diastolic ≥110mm of Hg (%) | | | | | | | | |
| Male | 2.2 | 2.5 | 2.0 | 2.0 | 2.7 | 2.2 | 1.1 | - |
| Female | 1.9 | 2.0 | 1.7 | 1.8 | 2.9 | 2.7 | 1.4 | 1.2 |
| Person | 2.0 | 2.2 | 1.8 | 1.9 | 2.8 | 2.5 | 1.2 | 0.9 |
| Iodine content in Household Salt (more than 15 ppm) (%) | 79.5 | 79.1 | 93.5 | 93.6 | 86.0 | 88.9 | 67.7 | 68.0 |

| Indicators | Mahasamund | | Dhamtari | | Kanker | | Bastar | | Dantewada | |
|--|------------|-------|----------|-------|--------|-------|--------|-------|-----------|-------|
| | Total | Rural | Total | Rural | Total | Rural | Total | Rural | Total | Rural |
| Nutritional status below 5 years | | | | | | | | | | |
| Below -2 SD Wasting (Weight for Height) (%) | | | | | | | | | | |
| Male | 35.1 | 33.6 | 33.0 | 35.0 | 19.4 | 19.6 | 6.6 | 6.6 | 23.6 | 20.4 |
| Female | 34.1 | 30.6 | 22.0 | 24.0 | 21.8 | 21.5 | 11.1 | 11.4 | 22.7 | 21.1 |
| Person | 34.7 | 32.2 | 26.8 | 28.8 | 20.6 | 20.5 | 8.5 | 8.6 | 23.1 | 20.7 |
| Below -3 SD Wasting (Weight for Height) (%) | | | | | | | | | | |
| Male | 14.6 | 12.8 | 9.0 | 6.9 | - | - | 2.8 | 2.5 | 5.0 | - |
| Female | 15.3 | 14.0 | 6.5 | 6.9 | - | - | - | - | 6.9 | 7.0 |
| Person | 14.9 | 13.3 | 7.6 | 6.9 | 3.2 | 3.3 | 3.8 | 3.5 | 5.9 | 5.9 |
| Below -2 SD Stunting (Height for Age) (%) | | | | | | | | | | |
| Male | 29.8 | 30.6 | 28.4 | 29.6 | 33.9 | 34.0 | 72.5 | 73.2 | 27.1 | 30.6 |
| Female | 35.7 | 38.9 | 33.5 | 33.1 | 25.0 | 24.6 | 46.4 | 45.7 | 23.5 | 25.0 |
| Person | 32.5 | 34.4 | 31.2 | 31.5 | 29.3 | 29.2 | 61.1 | 61.4 | 25.4 | 27.8 |
| Below -3SD Stunting (Height for Age) (%) | | | | | | | | | | |
| Male | 13.3 | 14.1 | 14.0 | 13.1 | 19.3 | 19.2 | 53.6 | 54.1 | 19.8 | 22.8 |
| Female | 17.9 | 19.5 | 11.4 | 10.2 | 11.3 | 11.2 | 30.9 | 31.1 | 13.5 | 14.7 |
| Person | 15.4 | 16.6 | 12.5 | 11.4 | 15.2 | 15.1 | 43.7 | 44.2 | 16.8 | 18.8 |
| Below -2 SD Underweight (Weight for Age) (%) | | | | | | | | | | |
| Male | 34.2 | 34.4 | 40.3 | 38.1 | 32.6 | 32.9 | 36.9 | 35.8 | 31.2 | 34.2 |
| Female | 33.2 | 34.3 | 35.8 | 36.6 | 24.6 | 24.8 | 26.6 | 26.4 | 20.8 | 21.6 |
| Person | 33.7 | 34.4 | 37.8 | 37.3 | 28.4 | 28.6 | 32.5 | 31.7 | 26.3 | 28.2 |
| Below -3 SD Underweight (Weight for Age) (%) | | | | | | | | | | |
| Male | 13.0 | 13.2 | 17.6 | 13.1 | 15.2 | 15.5 | 14.6 | 13.0 | 15.2 | 17.1 |
| Female | 16.4 | 17.8 | 13.9 | 13.4 | 11.4 | 11.2 | 9.4 | - | 7.2 | 7.9 |
| Person | 14.5 | 15.3 | 15.6 | 13.3 | 13.2 | 13.2 | 12.4 | 11.0 | 11.5 | 12.7 |
| Below -2 SD Undernourished (BMI for Age) (%) | | | | | | | | | | |
| Male | 39.7 | 38.6 | 31.7 | 35.3 | 19.3 | 19.6 | 7.1 | 7.2 | 32.1 | 30.3 |
| Female | 40.1 | 37.9 | 25.1 | 27.5 | 22.3 | 22.2 | 10.5 | 10.9 | 26.8 | 24.8 |
| Person | 39.9 | 38.3 | 28.1 | 30.9 | 20.8 | 21.0 | 8.5 | 8.8 | 29.7 | 27.7 |
| Below -3 SD Undernourished (BMI for Age) (%) | | | | | | | | | | |
| Male | 23.3 | 22.1 | 11.7 | 10.8 | 4.7 | 4.7 | 2.6 | 2.8 | 13.0 | 12.5 |
| Female | 22.0 | 21.2 | 6.4 | 6.3 | 4.0 | 4.1 | - | - | 10.3 | 9.8 |
| Person | 22.7 | 21.7 | 8.8 | 8.3 | 4.3 | 4.4 | 3.7 | 3.8 | 11.8 | 11.2 |
| Above 2 SD Overnourished (BMI for Age) (%) | | | | | | | | | | |
| Male | 2.8 | 3.1 | 9.6 | 6.0 | 3.6 | - | 20.5 | - | 8.0 | 8.4 |
| Female | 3.3 | 3.1 | 6.2 | - | - | - | 18.5 | 18.3 | 5.5 | 6.1 |
| Person | 3.1 | 3.1 | 7.7 | 4.5 | 4.9 | - | 19.6 | 19.2 | 6.8 | 7.3 |
| Above 3 SD Overnourished (BMI for Age) (%) | | | | | | | | | | |
| Male | 2.8 | 3.1 | - | - | - | - | 7.9 | - | 4.8 | 5.0 |
| Female | - | - | 2.2 | - | - | - | 6.3 | 5.6 | 3.7 | 4.1 |
| Person | 2.0 | 2.2 | 2.5 | - | 2.4 | - | 7.2 | 6.5 | 4.3 | 4.6 |
| Nutritional status 5-18 years | | | | | | | | | | |
| Below -2 SD Undernourished (BMI for Age) (%) | | | | | | | | | | |
| Male | 31.7 | 36.3 | 37.5 | 39.2 | 42.9 | 43.5 | 34.3 | 34.9 | 31.3 | 31.2 |
| Female | 24.5 | 27.2 | 27.5 | 29.9 | 30.7 | 31.2 | 27.6 | 28.5 | 24.7 | 24.9 |
| Person | 28.1 | 31.6 | 32.4 | 34.5 | 36.6 | 37.1 | 30.9 | 31.7 | 28.3 | 28.3 |
| Below -3 SD Undernourished (BMI for Age) (%) | | | | | | | | | | |
| Male | 13.1 | 15.0 | 16.5 | 17.5 | 18.3 | 18.5 | 18.3 | 18.4 | 16.6 | 16.8 |
| Female | 11.6 | 12.9 | 12.3 | 13.3 | 13.7 | 13.9 | 13.5 | 14.1 | 9.0 | 9.0 |
| Person | 12.4 | 13.9 | 14.4 | 15.4 | 15.9 | 16.1 | 15.8 | 16.2 | 13.1 | 13.3 |
| Above 2 SD Overnourished (BMI for Age) (%) | | | | | | | | | | |
| Male | - | - | 1.0 | 0.9 | - | - | 1.2 | 1.3 | 2.2 | 2.2 |
| Female | 1.0 | 0.9 | 0.7 | - | 1.7 | 1.7 | 1.1 | 1.1 | 1.3 | - |
| Person | 1.1 | 1.3 | 0.8 | 0.8 | 1.4 | 1.5 | 1.2 | 1.2 | 1.8 | 1.7 |
| Above 3 SD Overnourished (BMI for Age) (%) | | | | | | | | | | |
| Male | - | - | - | - | - | - | - | - | - | - |
| Female | 0.6 | 0.6 | - | - | - | - | 0.8 | 0.8 | - | - |
| Person | 0.3 | 0.3 | 0.4 | 0.3 | - | - | 0.7 | 0.8 | 1.2 | - |

| Indicators | Mahasamund | | Dhamtari | | Kanker | | Bastar | | Dantewada | |
|--|------------|-------|----------|-------|--------|-------|--------|-------|-----------|-------|
| | Total | Rural | Total | Rural | Total | Rural | Total | Rural | Total | Rural |
| Nutritional status 18 years and above | | | | | | | | | | |
| BMI ≥25.0 for age 18-59 year (%) | | | | | | | | | | |
| Male | 9.2 | 8.2 | 3.3 | 2.8 | 7.7 | 7.5 | 3.7 | 3.3 | 3.1 | 3.2 |
| Female | 9.7 | 10.1 | 7.6 | 6.2 | 6.3 | 5.6 | 2.1 | 2.1 | 1.4 | 1.3 |
| Person | 9.5 | 9.2 | 5.5 | 4.5 | 7.0 | 6.5 | 2.9 | 2.7 | 2.3 | 2.3 |
| BMI ≥25.0 for age 60 year and above (%) | | | | | | | | | | |
| Male | 11.8 | 11.3 | 4.1 | - | 5.6 | 5.3 | - | - | - | - |
| Female | 16.6 | 17.1 | 8.0 | 7.4 | 5.9 | 5.9 | - | - | 4.9 | 5.4 |
| Person | 14.2 | 14.4 | 6.0 | 5.0 | 5.7 | 5.6 | - | - | 3.3 | 3.7 |
| BMI ≥30.0 for age 18-59 year (%) | | | | | | | | | | |
| Male | 1.2 | 0.9 | 0.5 | 0.5 | 1.1 | 1.0 | - | - | 0.5 | 0.5 |
| Female | 1.8 | 1.8 | - | - | 1.0 | 0.8 | - | - | 0.5 | 0.5 |
| Person | 1.5 | 1.4 | 0.8 | 0.7 | 1.0 | 0.9 | 0.2 | 0.2 | 0.5 | 0.5 |
| BMI ≥30.0 for age 60 year and above (%) | | | | | | | | | | |
| Male | - | - | - | - | - | - | - | - | 0.0 | 0.0 |
| Female | 5.8 | 6.4 | - | - | - | - | - | - | 0.0 | 0.0 |
| Person | 3.3 | 3.9 | - | - | 1.2 | 1.0 | - | - | 0.0 | 0.0 |
| BMI <18.5 for age 18-59 year (%) | | | | | | | | | | |
| Male | 13.5 | 16.4 | 15.6 | 16.1 | 27.1 | 27.6 | 9.7 | 10.0 | 7.5 | 6.8 |
| Female | 19.3 | 22.3 | 19.8 | 21.2 | 35.6 | 36.3 | 12.5 | 12.8 | 25.0 | 24.9 |
| Person | 16.4 | 19.5 | 17.7 | 18.6 | 31.4 | 32.0 | 11.1 | 11.4 | 15.6 | 15.1 |
| BMI <18.5 for age 60 year and above (%) | | | | | | | | | | |
| Male | 16.5 | 20.7 | 28.6 | 28.2 | 42.0 | 43.0 | 12.7 | 11.4 | 46.0 | 46.6 |
| Female | 18.5 | 21.1 | 35.3 | 36.1 | 37.8 | 38.7 | 17.4 | 16.1 | 43.2 | 40.2 |
| Person | 17.5 | 20.9 | 31.9 | 32.2 | 39.9 | 40.8 | 14.8 | 13.5 | 44.7 | 43.6 |
| Anaemia status by Haemoglobin Level | | | | | | | | | | |
| Anaemia in the age group 6-59 months (%) | | | | | | | | | | |
| Male | 64.4 | 65.6 | 66.3 | 71.7 | 80.9 | 80.8 | 83.8 | 83.7 | 64.0 | 66.7 |
| Female | 56.3 | 54.4 | 70.1 | 71.8 | 67.5 | 67.4 | 76.5 | 76.2 | 71.1 | 70.1 |
| Person | 61.0 | 61.1 | 68.4 | 71.7 | 73.6 | 73.5 | 80.7 | 80.4 | 67.5 | 68.4 |
| Severe Anemia in the age group 6-59 months (%) | | | | | | | | | | |
| Male | - | - | - | - | - | - | 7.7 | 7.8 | - | - |
| Female | - | - | - | - | - | - | 13.4 | 13.6 | - | - |
| Person | - | - | 1.0 | 1.1 | - | - | 10.2 | 10.3 | - | - |
| Anaemia in the age group 5-9 years (%) | | | | | | | | | | |
| Male | 82.6 | 80.6 | 85.5 | 85.4 | 87.6 | 87.8 | 94.1 | 94.0 | 82.6 | 82.0 |
| Female | 84.0 | 83.8 | 82.5 | 84.6 | 84.3 | 84.5 | 91.9 | 91.7 | 84.5 | 83.1 |
| Person | 83.3 | 82.1 | 84.1 | 85.0 | 85.9 | 86.1 | 93.0 | 92.9 | 83.3 | 82.4 |
| Severe Anemia in the age group 5-9 years (%) | | | | | | | | | | |
| Male | 8.0 | 7.0 | 12.3 | 12.7 | - | - | 32.4 | 32.8 | 19.4 | 19.4 |
| Female | 8.9 | 5.4 | 9.1 | 9.1 | - | - | 33.3 | 33.7 | 13.2 | 13.6 |
| Person | 8.4 | 6.2 | 10.7 | 10.9 | - | - | 32.9 | 33.2 | 17.0 | 17.1 |
| Anaemia in the age group 10-17 years (%) | | | | | | | | | | |
| Male | 75.0 | 72.0 | 75.6 | 75.7 | 79.1 | 79.6 | 88.7 | 88.4 | 86.4 | 86.6 |
| Female | 77.3 | 75.9 | 76.4 | 77.7 | 82.6 | 82.8 | 90.6 | 90.6 | 79.3 | 78.8 |
| Person | 76.2 | 74.0 | 76.0 | 76.7 | 80.8 | 81.2 | 89.6 | 89.5 | 83.0 | 82.9 |
| Severe Anemia in the age group 10-17 years (%) | | | | | | | | | | |
| Male | 7.3 | 4.6 | 7.3 | 7.4 | 11.3 | 11.6 | 32.4 | 32.8 | 14.4 | 13.4 |
| Female | 8.0 | 5.2 | 8.2 | 7.8 | 8.9 | 9.0 | 27.5 | 28.0 | 9.5 | 9.4 |
| Person | 7.6 | 4.9 | 7.8 | 7.6 | 10.1 | 10.3 | 30.0 | 30.4 | 12.0 | 11.5 |
| Anaemia in the age group 18-59 years (%) | | | | | | | | | | |
| Male | 81.4 | 78.6 | 90.0 | 90.3 | 87.5 | 87.9 | 93.0 | 92.9 | 88.0 | 87.7 |
| Female | 78.8 | 77.4 | 79.1 | 79.7 | 82.4 | 82.8 | 89.5 | 89.3 | 78.6 | 77.5 |
| Severe Anemia in the age group 18-59 years (%) | | | | | | | | | | |
| Male | 5.1 | 3.6 | 10.7 | 11.4 | 10.1 | 10.3 | 31.1 | 31.6 | 9.8 | 9.8 |
| Female | 5.9 | 4.7 | 11.6 | 11.8 | 12.1 | 12.5 | 33.7 | 34.3 | 11.8 | 10.7 |
| Anaemia in the age group 60 years and above (%) | | | | | | | | | | |
| Male | 87.6 | 85.2 | 86.3 | 88.4 | 94.0 | 94.0 | 97.0 | 96.8 | 88.4 | 88.5 |
| Female | 82.5 | 80.7 | 84.2 | 84.8 | 86.4 | 86.3 | 92.1 | 91.5 | 79.6 | 79.6 |
| Severe Anemia in the age group 60 years and above (%) | | | | | | | | | | |
| Male | 6.8 | 5.9 | 14.1 | 16.2 | 10.6 | 10.6 | 38.9 | 39.9 | 16.0 | 14.9 |
| Female | 10.0 | 8.6 | 10.4 | 10.1 | 17.8 | 18.8 | 28.6 | 26.9 | 10.9 | 9.6 |

| Indicators | Mahasamund | | Dhamtari | | Kanker | | Bastar | | Dantewada | |
|--|------------|-------|----------|-------|--------|-------|--------|-------|-----------|-------|
| | Total | Rural | Total | Rural | Total | Rural | Total | Rural | Total | Rural |
| Blood Sugar Level (18 years and above) | | | | | | | | | | |
| Blood Sugar Level ≥110 mg/dl (%) | | | | | | | | | | |
| Male | 16.1 | 15.5 | 9.8 | 10.0 | 14.5 | 14.5 | 9.4 | 9.9 | 10.7 | 11.2 |
| Female | 10.8 | 10.8 | 7.5 | 7.6 | 11.6 | 11.4 | 4.8 | 5.2 | 8.3 | 8.6 |
| Person | 13.4 | 13.1 | 8.6 | 8.7 | 12.9 | 12.9 | 7.2 | 7.6 | 9.5 | 10.0 |
| Blood Sugar Level ≥130 mg/dl (%) | | | | | | | | | | |
| Male | 6.8 | 6.7 | 0.9 | 0.9 | 4.4 | 4.3 | 1.0 | 1.0 | 5.5 | 5.8 |
| Female | 4.9 | 5.0 | 0.8 | 0.8 | 4.2 | 4.0 | 0.5 | 0.6 | 3.8 | 3.9 |
| Person | 5.8 | 5.8 | 0.8 | 0.8 | 4.2 | 4.1 | 0.8 | 0.8 | 4.7 | 4.9 |
| Blood Sugar Level ≥150 mg/dl (%) | | | | | | | | | | |
| Male | 2.1 | 2.0 | 0.7 | 0.6 | 1.9 | 1.9 | 0.4 | 0.4 | 0.9 | 0.9 |
| Female | 1.6 | 1.6 | 0.5 | 0.5 | 2.1 | 2.0 | - | - | 0.8 | 0.8 |
| Person | 1.8 | 1.8 | 0.6 | 0.5 | 2.0 | 1.9 | 0.3 | 0.3 | 0.9 | 0.9 |
| Hypertension (18 years and above) | | | | | | | | | | |
| Above Normal Range | | | | | | | | | | |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%) | | | | | | | | | | |
| Male | 7.4 | 7.8 | 4.8 | 4.7 | 6.7 | 6.4 | 1.8 | 1.8 | 3.3 | 3.4 |
| Female | 8.4 | 8.6 | 4.7 | 4.7 | 6.0 | 5.8 | - | - | 2.2 | 2.4 |
| Person | 7.9 | 8.2 | 4.8 | 4.7 | 6.3 | 6.1 | 2.4 | 2.5 | 2.8 | 2.9 |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%) | | | | | | | | | | |
| Male | 7.7 | 7.0 | 4.4 | 4.1 | 7.5 | 7.6 | 3.3 | 3.4 | 2.0 | 2.1 |
| Female | 6.6 | 7.1 | 3.5 | 3.7 | 5.7 | 5.6 | 2.4 | 2.5 | 1.3 | 1.3 |
| Person | 7.2 | 7.0 | 4.0 | 3.9 | 6.6 | 6.6 | 2.9 | 2.9 | 1.6 | 1.7 |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%) | | | | | | | | | | |
| Male | 7.7 | 7.8 | 8.3 | 8.7 | 9.1 | 8.6 | 5.0 | 5.1 | 10.9 | 10.6 |
| Female | 6.4 | 6.5 | 6.7 | 6.9 | 6.7 | 6.4 | 4.4 | 4.6 | 9.0 | 8.6 |
| Person | 7.0 | 7.1 | 7.5 | 7.8 | 7.8 | 7.5 | 4.7 | 4.8 | 9.9 | 9.6 |
| Moderately high | | | | | | | | | | |
| Systolic ≥160mm of Hg & Diastolic ≥100mm of Hg (%) | | | | | | | | | | |
| Male | 2.4 | 2.8 | 0.6 | 0.5 | 1.9 | 1.8 | - | - | - | - |
| Female | 2.9 | 3.2 | 0.8 | 0.8 | 1.6 | 1.6 | - | - | - | - |
| Person | 2.7 | 3.0 | 0.7 | 0.6 | 1.8 | 1.7 | - | - | 0.1 | 0.1 |
| Systolic ≥160mm of Hg & Diastolic <100mm of Hg (%) | | | | | | | | | | |
| Male | 1.4 | - | 0.9 | 0.8 | 2.0 | 2.0 | - | - | 0.6 | - |
| Female | 2.0 | 2.1 | 1.1 | 1.0 | 1.7 | 1.5 | 0.7 | 0.7 | - | - |
| Person | 1.7 | 1.8 | 1.0 | 0.9 | 1.8 | 1.7 | 0.5 | 0.5 | 0.5 | 0.5 |
| Systolic <160mm of Hg & Diastolic ≥100mm of Hg (%) | | | | | | | | | | |
| Male | 2.6 | 2.5 | 1.7 | 1.5 | 2.9 | 2.6 | - | - | 4.6 | 4.8 |
| Female | 2.1 | 2.1 | 1.2 | 1.0 | 2.8 | 2.7 | 0.9 | 0.9 | 3.9 | 4.0 |
| Person | 2.4 | 2.3 | 1.4 | 1.3 | 2.9 | 2.7 | 0.7 | 0.7 | 4.2 | 4.4 |
| Very high | | | | | | | | | | |
| Systolic ≥180mm of Hg & Diastolic ≥110mm of Hg (%) | | | | | | | | | | |
| Male | 0.7 | 0.9 | - | - | 0.8 | 0.8 | - | - | - | - |
| Female | 0.5 | 0.6 | - | - | 0.5 | 0.4 | - | - | - | - |
| Person | 0.6 | 0.7 | 0.1 | 0.1 | 0.6 | 0.6 | - | - | - | - |
| Systolic ≥180mm of Hg & Diastolic <110mm of Hg (%) | | | | | | | | | | |
| Male | 0.8 | - | - | - | 0.5 | 0.5 | - | - | - | - |
| Female | 1.7 | 1.8 | 0.5 | 0.5 | 0.8 | 0.7 | - | - | - | - |
| Person | 1.2 | 1.3 | 0.3 | 0.3 | 0.7 | 0.6 | - | - | 0.2 | - |
| Systolic <180mm of Hg & Diastolic ≥110mm of Hg (%) | | | | | | | | | | |
| Male | 2.0 | 2.3 | 0.5 | 0.5 | 1.6 | 1.3 | - | - | 0.9 | 1.0 |
| Female | 1.3 | 1.2 | 0.4 | 0.4 | 0.9 | 0.8 | 0.4 | 0.4 | 0.6 | 0.6 |
| Person | 1.6 | 1.7 | 0.5 | 0.5 | 1.2 | 1.1 | 0.3 | 0.3 | 0.7 | 0.8 |
| Iodine content in Household Salt (more than 15 ppm) (%) | | | | | | | | | | |
| | 78.6 | 80.8 | 72.6 | 68.2 | 78.7 | 79.1 | 74.1 | 72.4 | 72.2 | 70.8 |

Note: "-" Indicate Inadequate Sample Size.

Anemia Cut-off.

Children (6-59 months) - Anemia Hb < 11 g/dL and Severe Anemia Hb < 5 g/dL

Male and Female aged 5-17 years- Anemia Hb < 12g/dL and Severe Anemia Hb < 7g/dL

Men (18 years and above)- Anemia Hb < 13 g/dL and Severe Anemia Hb < 7g/dL

Non-Pregnant Women (18 years and above)- Anemia Hb < 12g/dL and Severe Anemia Hb < 7g/dL