

| Indicators  | Madhya Pradesh |                |      |             |             |             |                |      |             |             |
|---|----------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total          |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size    | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |                |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |                |                |      |             |             |             |                |      |             |             |
| Person  | 15884          | 0.7            | 3.9  | 16.0        | 18.6        | 11685       | 0.8            | 4.2  | 16.9        | 20.0        |
| Below -2 SD Stunting (Height for Age) (%)               |                |                |      |             |             |             |                |      |             |             |
| Person  | 15934          | 1.2            | 2.4  | 49.1        | 53.9        | 11732       | 1.3            | 2.7  | 47.0        | 52.3        |
| Below -2 SD Underweight (Weight for Age) (%)            |                |                |      |             |             |             |                |      |             |             |
| Person  | 16887          | 1.0            | 2.5  | 38.7        | 42.6        | 12432       | 1.1            | 2.8  | 37.5        | 41.8        |
| Below -2 SD Undernourished (BMI for Age) (%)            |                |                |      |             |             |             |                |      |             |             |
| Person  | 17239          | 0.8            | 3.8  | 18.3        | 21.3        | 12673       | 0.9            | 4.3  | 19.1        | 22.6        |
| Above 2 SD Overnourished (BMI for Age) (%)              |                |                |      |             |             |             |                |      |             |             |
| Person  | 17239          | 0.6            | 5.3  | 10.3        | 12.7        | 12673       | 0.7            | 6.2  | 10.4        | 13.3        |
| <b>Nutritional status 5-18 years</b>                    |                |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |                |                |      |             |             |             |                |      |             |             |
| Person  | 61615          | 0.8            | 2.6  | 29.6        | 32.8        | 45163       | 1.1            | 3.3  | 30.7        | 35.0        |
| Above 2 SD Overnourished (BMI for Age) (%)              |                |                |      |             |             |             |                |      |             |             |
| Person  | 61615          | 0.1            | 6.4  | 1.5         | 1.9         | 45163       | 0.1            | 6.0  | 1.4         | 1.8         |
| <b>Nutritional status 18 years and above</b>            |                |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |                |                |      |             |             |             |                |      |             |             |
| Person  | 131279         | 0.4            | 5.8  | 5.8         | 7.6         | 89396       | 0.2            | 5.2  | 4.0         | 4.9         |
| BMI ≥25.0 for age 60 year and above (%)                 |                |                |      |             |             |             |                |      |             |             |
| Person  | 20683          | 0.6            | 6.6  | 8.3         | 11.0        | 14679       | 0.3            | 5.6  | 5.5         | 6.8         |
| BMI <18.5 for age 18-59 year (%)                        |                |                |      |             |             |             |                |      |             |             |
| Person  | 131279         | 0.9            | 3.8  | 21.2        | 25.3        | 89396       | 1.2            | 4.9  | 21.8        | 27.2        |
| BMI <18.5 for age 60 year and above (%)                 |                |                |      |             |             |             |                |      |             |             |
| Person  | 20683          | 1.1            | 4.3  | 23.4        | 28.4        | 14679       | 1.4            | 4.7  | 26.2        | 32.6        |
| <b>Anaemia status by Haemoglobin Level</b>              |                |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |                |                |      |             |             |             |                |      |             |             |
| Person  | 7981           | 1.0            | 1.3  | 74.3        | 78.2        | 6090        | 1.1            | 1.5  | 74.3        | 78.7        |
| Anaemia in the age group 5-9 years (%)                  |                |                |      |             |             |             |                |      |             |             |
| Person  | 14052          | 0.8            | 0.9  | 83.3        | 86.4        | 10590       | 0.9            | 1.1  | 83.4        | 87.0        |
| Anaemia in the age group 10-17 years (%)                |                |                |      |             |             |             |                |      |             |             |
| Person  | 30221          | 0.8            | 0.9  | 80.8        | 83.9        | 22385       | 0.9            | 1.1  | 80.6        | 84.1        |
| Anaemia in the age group 18-59 years (%)                |                |                |      |             |             |             |                |      |             |             |
| Male  | 55640          | 0.8            | 1.0  | 82.9        | 86.1        | 39059       | 1.0            | 1.2  | 82.5        | 86.4        |
| Female  | 55317          | 0.8            | 0.9  | 82.3        | 85.3        | 38119       | 0.9            | 1.1  | 81.7        | 85.3        |
| Anaemia in the age group 60 years and above (%)         |                |                |      |             |             |             |                |      |             |             |
| Male  | 9201           | 0.8            | 0.9  | 86.1        | 89.3        | 6628        | 1.0            | 1.2  | 84.8        | 88.8        |
| Female  | 8427           | 1.0            | 1.2  | 82.6        | 86.6        | 6074        | 1.2            | 1.4  | 81.7        | 86.5        |
| <b>Blood Sugar Level (18 years and above)</b>           |                |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |                |                |      |             |             |             |                |      |             |             |
| Person  | 141287         | 0.3            | 3.6  | 8.2         | 9.5         | 97007       | 0.4            | 4.8  | 7.3         | 8.8         |
| <b>Hypertension (18 years and above)</b>                |                |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |                |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |                |                |      |             |             |             |                |      |             |             |
| Person  | 151240         | 0.3            | 3.1  | 10.0        | 11.3        | 103066      | 0.4            | 4.1  | 8.7         | 10.2        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |                |                |      |             |             |             |                |      |             |             |
| Person  | 151240         | 0.3            | 5.0  | 4.6         | 5.6         | 103066      | 0.3            | 5.6  | 4.2         | 5.2         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |                |                |      |             |             |             |                |      |             |             |
| Person  | 151240         | 0.3            | 4.8  | 5.6         | 6.8         | 103066      | 0.4            | 5.8  | 6.2         | 7.8         |
| Iodine content in Household Salt (more than 15 ppm) (%) | 58094          | 1.8            | 2.5  | 68.8        | 76.1        | 41375       | 2.2            | 3.3  | 61.7        | 70.4        |

| Indicators  | Sheopur     |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 588         | 6.3            | 34.7 | 7.9         | 36.9        | 504         | 7.2            | 36.9 | 8.0         | 40.8        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 604         | 6.2            | 10.8 | 43.3        | 70.6        | 520         | 6.8            | 11.9 | 42.0        | 72.0        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 612         | 9.3            | 19.3 | 28.5        | 68.3        | 528         | 10.5           | 21.7 | 26.6        | 70.9        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 615         | 3.8            | 30.0 | 6.3         | 24.3        | 530         | 4.3            | 32.1 | 6.3         | 26.5        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 615         | 4.0            | 47.2 | 2.8         | 23.1        | -           | -              | -    | -           | -           |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1679        | 1.5            | 6.6  | 20.1        | 27.0        | 1257        | 1.8            | 7.4  | 21.0        | 29.3        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | -           | -              | -    | -           | -           | -           | -              | -    | -           | -           |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2699        | 0.8            | 27.4 | 1.6         | 5.5         | 2160        | 0.9            | 33.6 | 1.2         | 5.6         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 431         | 1.8            | 31.0 | 2.9         | 11.7        | 338         | 1.5            | 37.1 | 1.7         | 9.3         |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2699        | 2.8            | 10.0 | 22.2        | 34.9        | 2160        | 3.3            | 11.8 | 21.0        | 35.7        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 431         | 4.4            | 18.2 | 15.6        | 35.4        | 338         | 5.2            | 20.9 | 15.0        | 38.5        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 348         | 10.8           | 16.8 | 38.3        | 83.7        | 282         | 13.2           | 22.0 | 30.1        | 83.9        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 433         | 10.4           | 15.5 | 41.3        | 85.7        | 364         | 12.2           | 18.9 | 35.3        | 85.8        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 675         | 9.2            | 15.1 | 39.6        | 79.3        | 535         | 11.5           | 20.6 | 30.6        | 78.4        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1123        | 3.9            | 7.0  | 47.2        | 64.8        | 938         | 4.3            | 8.2  | 42.1        | 61.2        |
| Female  | 1069        | 7.2            | 11.4 | 45.9        | 77.6        | 858         | 8.7            | 15.1 | 38.0        | 75.6        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 214         | 6.5            | 9.8  | 50.5        | 79.2        | 171         | 7.5            | 12.0 | 44.6        | 77.3        |
| Female  | 166         | 10.4           | 17.6 | 35.3        | 79.4        | 127         | 11.7           | 23.2 | 26.1        | 74.6        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2758        | 2.4            | 21.8 | 6.8         | 18.1        | 2279        | 2.8            | 26.1 | 5.9         | 19.1        |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3077        | 2.4            | 16.7 | 9.7         | 20.6        | 2456        | 2.8            | 18.5 | 9.8         | 22.7        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3077        | 0.7            | 38.5 | 0.7         | 4.1         | -           | -              | -    | -           | -           |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3077        | 0.5            | 27.5 | 0.9         | 3.1         | 2456        | 0.6            | 30.3 | 0.9         | 3.7         |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1416        | 13.7           | 20.0 | 34.0        | 90.1        | 1179        | 15.5           | 23.0 | 29.5        | 91.0        |

| Indicators  | Morena      |                |            |             |             |             |                |            |             |             |
|---|-------------|----------------|------------|-------------|-------------|-------------|----------------|------------|-------------|-------------|
|   | Total       |                |            |             |             | Rural       |                |            |             |             |
|   | Sample Size | Standard Error | PRSE       | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE       | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |            |             |             |             |                |            |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |            |             |             |             |                |            |             |             |
| Person  | 423         | 2.6            | 16.4       | 10.9        | 22.9        | 339         | 2.0            | 13.1       | 11.5        | 20.7        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |            |             |             |             |                |            |             |             |
| Person  | 446         | 5.4            | 9.8        | 42.8        | 66.6        | 349         | 6.2            | 11.5       | 39.7        | 66.9        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |            |             |             |             |                |            |             |             |
| Person  | 448         | 9.3            | 28.7       | 15.5        | 55.5        | 351         | 6.5            | 25.8       | 13.4        | 42.5        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |            |             |             |             |                |            |             |             |
| Person  | 448         | 2.2            | 10.6       | 16.2        | 26.1        | 351         | 2.5            | 13.7       | 13.3        | 24.6        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |            |             |             |             |                |            |             |             |
| Person  | 448         | 4.6            | 36.8       | 5.2         | 27.1        | 351         | 5.4            | 39.8       | 5.2         | 30.6        |
| <b>Nutritional status 5-18 years</b>                    |             |                |            |             |             |             |                |            |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |            |             |             |             |                |            |             |             |
| Person  | 1162        | 1.3            | 6.5        | 16.8        | 22.5        | 841         | 1.7            | 9.0        | 15.2        | 22.8        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |            |             |             |             |                |            |             |             |
| Person  | -           | -              | -          | -           | -           | -           | -              | -          | -           | -           |
| <b>Nutritional status 18 years and above</b>            |             |                |            |             |             |             |                |            |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |            |             |             |             |                |            |             |             |
| Person  | 2136        | 0.2            | 8.8        | 2.3         | 3.5         | 1508        | 0.3            | 11.4       | 2.0         | 3.4         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |            |             |             |             |                |            |             |             |
| Person  | 343         | 0.8            | 17.7       | 3.0         | 6.8         | 262         | 0.7            | 23.1       | 1.9         | 5.4         |
| BMI <18.5 for age 18-59 year (%)                        |             |                |            |             |             |             |                |            |             |             |
| Person  | 2136        | 1.1            | 6.7        | 13.8        | 18.7        | 1508        | 1.4            | 8.2        | 14.0        | 20.3        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |            |             |             |             |                |            |             |             |
| Person  | 343         | 1.8            | 18.2       | 6.4         | 14.6        | 262         | 1.4            | 14.8       | 6.7         | 13.0        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |            |             |             |             |                |            |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |            |             |             |             |                |            |             |             |
| Person  | 68          | 4.4            | 6.2        | 59.6        | 79.2        | 50          | 5.2            | 7.1        | 60.4        | 83.8        |
| Anaemia in the age group 5-9 years (%)                  |             |                |            |             |             |             |                |            |             |             |
| Person  | 185         | 3.8            | 5.0        | 65.4        | 82.4        | 123         | 5.0            | 6.6        | 62.4        | 84.7        |
| Anaemia in the age group 10-17 years (%)                |             |                |            |             |             |             |                |            |             |             |
| Person  | 541         | 2.8            | 3.7        | 68.6        | 81.0        | 382         | 2.8            | 3.6        | 70.0        | 82.6        |
| Anaemia in the age group 18-59 years (%)                |             |                |            |             |             |             |                |            |             |             |
| Male  | 1029        | 2.2            | 2.6        | 79.9        | 89.9        | 719         | 3.1            | 3.7        | 76.0        | 90.0        |
| Female  | 961         | 4.3            | 5.3        | 68.9        | 88.2        | 706         | 5.8            | 7.3        | 63.5        | 89.8        |
| Anaemia in the age group 60 years and above (%)         |             |                |            |             |             |             |                |            |             |             |
| Male  | 177         | 3.6            | 4.5        | 71.1        | 87.5        | 134         | 4.1            | 5.1        | 70.6        | 89.4        |
| Female  | 137         | 3.0            | 3.6        | 77.1        | 91.0        | 106         | 3.4            | 4.2        | 73.6        | 89.3        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |            |             |             |             |                |            |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |            |             |             |             |                |            |             |             |
| Person  | 2412        | 2.8            | 26.6       | 5.6         | 18.6        | 1716        | 4.1            | 31.5       | 6.2         | 25.4        |
| <b>Hypertension (18 years and above)</b>                |             |                |            |             |             |             |                |            |             |             |
| <b>Above Normal Range</b>                               |             |                |            |             |             |             |                |            |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |            |             |             |             |                |            |             |             |
| Person  | 2453        | 1.2            | 9.4        | 10.4        | 15.9        | 1753        | 1.6            | 11.4       | 11.2        | 18.7        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |            |             |             |             |                |            |             |             |
| Person  | -           | -              | -          | -           | -           | -           | -              | -          | -           | -           |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |            |             |             |             |                |            |             |             |
| Person  | 2453        | 1.2            | 26.4       | 2.4         | 8.0         | 1753        | 1.4            | 31.6       | 2.1         | 8.7         |
| Iodine content in Household Salt (more than 15 ppm) (%) | <b>1027</b> | <b>4.8</b>     | <b>6.7</b> | <b>59.5</b> | <b>80.8</b> | <b>738</b>  | <b>5.6</b>     | <b>8.0</b> | <b>56.6</b> | <b>81.6</b> |

| Indicators  | Bhind       |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 582         | 1.4            | 14.4 | 7.2         | 13.8        | 409         | 1.5            | 15.2 | 6.9         | 13.7        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 608         | 6.7            | 10.5 | 48.0        | 77.7        | 428         | 10.1           | 16.6 | 37.2        | 80.1        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 612         | 5.5            | 10.8 | 38.5        | 62.8        | 432         | 8.0            | 16.5 | 31.6        | 66.4        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 613         | 3.0            | 23.1 | 7.6         | 21.5        | 433         | 3.8            | 28.4 | 6.9         | 24.6        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 613         | 2.6            | 27.2 | 5.0         | 17.1        | 433         | 2.8            | 39.4 | 2.8         | 16.5        |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1344        | 4.1            | 16.4 | 17.0        | 35.7        | 690         | 4.6            | 16.0 | 19.5        | 40.2        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 1344        | 0.3            | 29.8 | 0.5         | 1.9         | 690         | 0.6            | 43.4 | 0.5         | 3.6         |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3186        | 0.4            | 15.3 | 2.0         | 3.9         | 1546        | 0.6            | 20.9 | 1.7         | 4.4         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 598         | 0.7            | 16.9 | 2.7         | 5.8         | 354         | 1.1            | 27.1 | 2.3         | 7.7         |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3186        | 3.2            | 16.6 | 13.1        | 27.7        | 1546        | 3.3            | 11.1 | 23.3        | 38.3        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 598         | 2.8            | 15.9 | 12.2        | 25.0        | 354         | 3.5            | 13.4 | 19.2        | 35.2        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 133         | 4.8            | 7.5  | 52.2        | 73.5        | 60          | 7.2            | 10.1 | 53.1        | 85.0        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 255         | 6.3            | 8.6  | 56.5        | 84.3        | 104         | 2.9            | 3.2  | 80.0        | 93.2        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 643         | 4.0            | 5.6  | 62.8        | 81.0        | 288         | 4.3            | 5.3  | 69.5        | 88.9        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1364        | 3.1            | 3.6  | 76.5        | 90.5        | 642         | 3.6            | 3.9  | 78.9        | 96.0        |
| Female  | 1277        | 4.0            | 5.5  | 63.4        | 81.5        | 620         | 3.4            | 4.1  | 75.3        | 91.0        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 283         | 3.9            | 4.7  | 72.5        | 90.2        | 168         | 4.4            | 4.9  | 74.7        | 96.4        |
| Female  | 221         | 4.8            | 6.3  | 63.8        | 85.6        | 129         | 7.1            | 8.5  | 61.3        | 94.1        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3451        | 1.4            | 13.4 | 7.9         | 14.4        | 1637        | 2.8            | 31.0 | 4.4         | 17.7        |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3986        | 1.1            | 7.3  | 12.3        | 17.2        | 2070        | 2.1            | 14.5 | 10.2        | 19.6        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3986        | 1.1            | 23.9 | 2.8         | 8.1         | 2070        | 2.3            | 40.5 | 2.2         | 13.6        |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3986        | 1.8            | 30.5 | 2.9         | 11.6        | 2070        | 2.3            | 27.8 | 4.4         | 15.3        |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1425        | 6.9            | 8.1  | 62.6        | 95.0        | 753         | 12.5           | 17.4 | 38.7        | 91.4        |

| Indicators  | Gwalior     |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 505         | 2.9            | 13.8 | 15.3        | 28.6        | 222         | 4.7            | 19.2 | 15.6        | 36.9        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 522         | 4.4            | 9.5  | 36.8        | 56.6        | 235         | 2.1            | 6.1  | 30.2        | 39.7        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 531         | 2.7            | 7.4  | 31.1        | 43.5        | 237         | 2.3            | 7.5  | 26.1        | 36.6        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 542         | 3.6            | 13.2 | 19.8        | 35.9        | 239         | 4.1            | 13.5 | 21.7        | 39.9        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 542         | 2.0            | 31.6 | 3.1         | 12.8        | 239         | 0.3            | 6.0  | 4.3         | 5.7         |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1326        | 3.2            | 10.2 | 24.8        | 39.3        | 450         | 3.9            | 12.7 | 22.5        | 39.9        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | -           | -              | -    | -           | -           | -           | -              | -    | -           | -           |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3544        | 0.9            | 15.0 | 4.4         | 8.7         | 1265        | 1.1            | 20.8 | 3.2         | 8.2         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 572         | 3.6            | 35.1 | 4.5         | 21.6        | 231         | 1.7            | 39.5 | 1.7         | 10.1        |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3544        | 2.3            | 8.0  | 24.0        | 34.4        | 1265        | 3.5            | 10.9 | 24.4        | 39.9        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 572         | 3.7            | 16.1 | 15.8        | 32.6        | 231         | 3.1            | 13.0 | 17.5        | 31.5        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 144         | 4.7            | 6.4  | 61.1        | 82.0        | 53          | 1.5            | 2.0  | 69.8        | 76.5        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 286         | 5.4            | 6.9  | 63.9        | 88.2        | 91          | 12.1           | 16.3 | 40.7        | 92.2        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 610         | 3.8            | 4.8  | 68.6        | 85.7        | 189         | 2.4            | 3.0  | 75.6        | 86.6        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1115        | 2.8            | 3.4  | 75.9        | 88.8        | 349         | 5.7            | 6.7  | 67.0        | 93.3        |
| Female  | 1348        | 3.2            | 3.9  | 73.4        | 87.8        | 408         | 4.4            | 5.3  | 71.1        | 91.4        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 214         | 3.0            | 3.4  | 79.9        | 94.0        | 76          | 6.5            | 7.5  | 64.5        | 95.5        |
| Female  | 204         | 4.1            | 4.7  | 74.7        | 93.6        | 78          | 3.3            | 3.5  | 80.9        | 97.1        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3600        | 3.4            | 30.8 | 5.4         | 21.5        | 1293        | 0.6            | 12.1 | 3.9         | 6.7         |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 4289        | 1.5            | 10.9 | 10.8        | 17.7        | 1527        | 0.8            | 8.2  | 8.2         | 11.9        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 4289        | 2.3            | 29.7 | 3.9         | 15.0        | 1527        | 1.1            | 19.5 | 3.6         | 8.7         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 4289        | 0.3            | 6.4  | 4.5         | 6.0         | 1527        | 0.7            | 16.1 | 3.0         | 6.2         |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1421        | 2.2            | 2.3  | 87.2        | 98.5        | 544         | 8.6            | 9.5  | 51.8        | 98.6        |

| Indicators  | Data        |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 445         | 4.3            | 13.1 | 24.1        | 43.5        | 239         | 6.4            | 19.5 | 20.2        | 48.5        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 466         | 3.6            | 6.2  | 50.3        | 66.5        | 251         | 4.6            | 7.6  | 49.6        | 69.9        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 472         | 6.2            | 9.5  | 50.5        | 77.9        | 255         | 9.7            | 15.2 | 40.6        | 81.9        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 473         | 3.1            | 12.7 | 18.0        | 31.9        | 256         | 3.3            | 13.5 | 17.8        | 32.8        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 473         | 0.9            | 24.0 | 2.3         | 6.7         | 256         | 1.3            | 30.9 | 2.0         | 8.2         |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1128        | 2.2            | 12.4 | 13.2        | 23.2        | 694         | 3.0            | 17.2 | 11.8        | 25.6        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 1128        | 0.8            | 47.2 | 0.6         | 4.8         | 694         | 1.1            | 49.3 | 0.8         | 6.9         |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2379        | 1.6            | 35.5 | 1.9         | 9.6         | 1364        | 2.4            | 42.9 | 2.1         | 14.2        |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 435         | 3.1            | 45.6 | 2.3         | 17.8        | -           | -              | -    | -           | -           |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2379        | 3.3            | 20.1 | 10.3        | 25.4        | 1364        | 4.3            | 27.4 | 8.3         | 28.2        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 435         | 2.8            | 22.1 | 7.6         | 20.5        | 265         | 3.2            | 35.2 | 4.0         | 19.5        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 112         | 3.1            | 4.7  | 57.9        | 71.7        | 74          | 4.2            | 6.7  | 52.3        | 71.1        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 265         | 4.2            | 5.4  | 66.8        | 85.6        | 159         | 3.6            | 4.5  | 70.0        | 86.3        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 620         | 5.0            | 7.3  | 56.9        | 79.3        | 347         | 6.0            | 8.5  | 55.6        | 82.5        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1006        | 3.5            | 4.3  | 71.1        | 86.7        | 529         | 3.9            | 4.8  | 70.6        | 88.3        |
| Female  | 950         | 5.5            | 7.3  | 60.8        | 85.2        | 486         | 4.3            | 5.7  | 64.8        | 84.3        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 189         | 3.2            | 3.9  | 72.8        | 87.2        | 104         | 4.2            | 5.2  | 68.4        | 87.2        |
| Female  | 163         | 5.0            | 6.9  | 60.4        | 82.9        | 88          | 7.1            | 9.9  | 53.1        | 84.3        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2651        | 2.1            | 18.4 | 7.4         | 16.8        | 1519        | 2.4            | 29.9 | 4.1         | 15.7        |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 2760        | 2.5            | 18.7 | 8.7         | 20.2        | 1605        | 1.4            | 13.6 | 7.4         | 13.6        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 2760        | 0.6            | 28.2 | 1.2         | 4.1         | 1605        | 0.6            | 33.1 | 0.9         | 3.8         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 2760        | 1.7            | 32.9 | 2.5         | 10.8        | 1605        | 1.9            | 41.2 | 1.8         | 11.3        |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1211        | 8.7            | 13.0 | 45.4        | 83.3        | 701         | 14.0           | 25.8 | 24.9        | 80.8        |

| Indicators  | Shivpuri    |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 595         | 1.6            | 14.0 | 8.2         | 15.4        | 517         | 1.8            | 15.7 | 8.1         | 16.4        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 600         | 2.5            | 4.2  | 53.6        | 64.7        | 522         | 1.2            | 2.0  | 56.1        | 61.3        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 607         | 6.9            | 18.2 | 23.9        | 54.3        | 527         | 7.6            | 20.5 | 21.9        | 55.0        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 628         | 3.6            | 23.1 | 9.1         | 25.8        | 547         | 4.1            | 25.3 | 9.0         | 28.0        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 628         | 1.1            | 20.0 | 3.5         | 8.5         | 547         | 0.8            | 20.1 | 2.6         | 6.5         |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 2133        | 1.4            | 3.6  | 36.3        | 42.7        | 1850        | 1.2            | 2.8  | 38.9        | 44.2        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | -           | -              | -    | -           | -           | -           | -              | -    | -           | -           |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3635        | 0.4            | 13.9 | 1.9         | 3.6         | 3043        | 0.4            | 22.5 | 1.2         | 3.3         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 640         | 1.6            | 28.8 | 2.8         | 10.3        | 558         | 1.8            | 39.2 | 1.8         | 10.7        |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3635        | 2.6            | 6.1  | 36.5        | 47.9        | 3043        | 3.0            | 7.0  | 35.6        | 48.9        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 640         | 2.1            | 5.4  | 35.1        | 44.8        | 558         | 2.5            | 5.9  | 36.4        | 47.6        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 316         | 4.5            | 6.0  | 63.1        | 83.3        | 279         | 4.9            | 6.5  | 62.8        | 85.0        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 400         | 4.3            | 4.8  | 75.3        | 97.1        | 337         | 5.3            | 5.8  | 70.4        | 97.4        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 964         | 2.3            | 2.6  | 81.1        | 91.6        | 823         | 2.3            | 2.7  | 80.4        | 91.0        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1238        | 1.5            | 1.7  | 85.5        | 92.5        | 1054        | 1.6            | 1.8  | 84.7        | 91.9        |
| Female  | 1330        | 1.8            | 2.0  | 82.5        | 90.5        | 1103        | 1.6            | 1.9  | 82.5        | 90.0        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 242         | 2.4            | 2.6  | 85.5        | 97.8        | 214         | 2.7            | 2.9  | 83.8        | 97.7        |
| Female  | 208         | 6.2            | 7.2  | 65.5        | 94.3        | 174         | 7.2            | 8.8  | 60.1        | 93.3        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3676        | 0.6            | 5.1  | 10.6        | 13.3        | 3087        | 0.7            | 6.4  | 10.0        | 13.4        |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 4307        | 1.1            | 7.7  | 12.3        | 17.5        | 3633        | 1.3            | 8.9  | 11.7        | 17.4        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 4307        | 1.2            | 20.3 | 3.8         | 9.5         | 3633        | 1.0            | 17.5 | 3.7         | 8.1         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 4307        | 0.8            | 15.3 | 3.9         | 7.7         | 3633        | 1.0            | 17.6 | 3.8         | 8.4         |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1539        | 15.9           | 27.6 | 23.8        | 85.8        | 1330        | 19.0           | 37.7 | 15.4        | 84.9        |

| Indicators  | Guna        |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 455         | 4.5            | 35.7 | 5.4         | 26.8        | 348         | 5.2            | 48.3 | 3.4         | 29.0        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 491         | 5.3            | 12.1 | 32.2        | 55.7        | 376         | 6.5            | 16.3 | 26.2        | 54.7        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 495         | 5.6            | 16.4 | 22.6        | 47.4        | 379         | 4.2            | 14.9 | 19.8        | 38.7        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 496         | 5.7            | 32.3 | 8.2         | 34.5        | 380         | 6.6            | 41.6 | 5.8         | 36.8        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 496         | 1.8            | 19.6 | 5.9         | 14.3        | 380         | 1.9            | 18.8 | 6.6         | 15.4        |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1213        | 4.8            | 16.5 | 19.4        | 40.7        | 933         | 5.8            | 19.0 | 19.1        | 44.7        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 1213        | 0.5            | 43.6 | 0.4         | 2.8         | 933         | 0.2            | 37.6 | 0.2         | 1.3         |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2241        | 0.5            | 15.5 | 2.4         | 4.8         | 1621        | 0.5            | 16.7 | 1.9         | 4.0         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 356         | 2.4            | 37.3 | 2.7         | 14.5        | -           | -              | -    | -           | -           |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2241        | 2.7            | 15.2 | 12.4        | 24.6        | 1621        | 2.6            | 13.5 | 14.2        | 26.0        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 356         | 3.5            | 25.1 | 7.8         | 23.9        | 263         | 4.7            | 30.3 | 7.5         | 29.0        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 185         | 9.5            | 16.4 | 36.3        | 76.8        | 156         | 10.7           | 18.3 | 34.1        | 79.1        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 241         | 4.5            | 5.7  | 66.8        | 86.9        | 207         | 4.6            | 5.7  | 67.6        | 88.2        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 456         | 3.3            | 4.0  | 74.5        | 89.5        | 376         | 3.2            | 3.8  | 74.8        | 89.1        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 922         | 3.0            | 3.4  | 80.2        | 94.4        | 747         | 3.3            | 3.7  | 79.1        | 94.7        |
| Female  | 791         | 3.7            | 4.6  | 70.6        | 87.4        | 671         | 3.6            | 4.5  | 71.7        | 88.1        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 152         | 4.4            | 5.0  | 73.9        | 94.8        | 126         | 4.9            | 5.5  | 72.4        | 96.3        |
| Female  | 127         | 6.7            | 9.2  | 55.3        | 84.9        | 96          | 7.2            | 9.5  | 56.4        | 88.4        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2540        | 1.4            | 10.5 | 10.4        | 16.7        | 1854        | 1.5            | 11.0 | 10.6        | 17.5        |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 2570        | 1.0            | 6.3  | 13.6        | 18.0        | 1865        | 1.1            | 7.6  | 12.7        | 17.9        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 2570        | 0.3            | 17.5 | 1.0         | 2.3         | 1865        | 0.4            | 23.6 | 0.9         | 2.6         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 2570        | 1.4            | 21.9 | 3.8         | 10.3        | 1865        | 1.7            | 30.8 | 2.7         | 10.7        |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1088        | 14.0           | 28.2 | 21.7        | 77.8        | 783         | 15.5           | 36.9 | 14.7        | 75.5        |



| Indicators  | Tikamgarh   |                |            |             |             |             |                |            |             |             |
|---|-------------|----------------|------------|-------------|-------------|-------------|----------------|------------|-------------|-------------|
|   | Total       |                |            |             |             | Rural       |                |            |             |             |
|   | Sample Size | Standard Error | PRSE       | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE       | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |            |             |             |             |                |            |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |            |             |             |             |                |            |             |             |
| Person  | 184         | 2.8            | 17.1       | 11.1        | 24.0        | 152         | 2.3            | 11.6       | 14.8        | 25.0        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |            |             |             |             |                |            |             |             |
| Person  | 177         | 3.3            | 4.9        | 59.8        | 74.8        | 149         | 3.8            | 5.6        | 59.4        | 76.5        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |            |             |             |             |                |            |             |             |
| Person  | 198         | 2.8            | 4.8        | 52.2        | 64.9        | 165         | 3.3            | 5.4        | 52.2        | 66.8        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |            |             |             |             |                |            |             |             |
| Person  | 198         | 2.7            | 13.3       | 14.6        | 26.7        | 165         | 3.1            | 13.8       | 16.5        | 30.6        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |            |             |             |             |                |            |             |             |
| Person  | 198         | 2.6            | 14.3       | 13.3        | 25.3        | 165         | 2.9            | 16.3       | 12.2        | 25.4        |
| <b>Nutritional status 5-18 years</b>                    |             |                |            |             |             |             |                |            |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |            |             |             |             |                |            |             |             |
| Person  | 1283        | 2.2            | 6.0        | 32.6        | 42.7        | 1067        | 2.3            | 6.1        | 33.5        | 44.1        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |            |             |             |             |                |            |             |             |
| Person  | 1283        | 0.4            | 23.5       | 1.1         | 3.1         | 1067        | 0.4            | 27.9       | 0.8         | 2.9         |
| <b>Nutritional status 18 years and above</b>            |             |                |            |             |             |             |                |            |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |            |             |             |             |                |            |             |             |
| Person  | 2702        | 1.8            | 34.7       | 2.3         | 11.1        | 2139        | 0.7            | 27.9       | 1.2         | 4.6         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |            |             |             |             |                |            |             |             |
| Person  | 493         | 1.7            | 45.0       | 1.3         | 10.2        | 415         | 0.9            | 46.9       | 0.6         | 5.5         |
| BMI <18.5 for age 18-59 year (%)                        |             |                |            |             |             |             |                |            |             |             |
| Person  | 2702        | 3.3            | 12.4       | 20.1        | 35.0        | 2139        | 3.5            | 11.9       | 22.4        | 38.2        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |            |             |             |             |                |            |             |             |
| Person  | 493         | 6.0            | 13.9       | 30.7        | 57.2        | 415         | 6.8            | 14.7       | 31.4        | 61.3        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |            |             |             |             |                |            |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |            |             |             |             |                |            |             |             |
| Person  | 194         | 3.1            | 3.7        | 76.4        | 90.7        | 156         | 3.1            | 3.6        | 76.1        | 90.1        |
| Anaemia in the age group 5-9 years (%)                  |             |                |            |             |             |             |                |            |             |             |
| Person  | 351         | 0.8            | 0.8        | 92.6        | 96.1        | 298         | 0.7            | 0.7        | 94.0        | 97.1        |
| Anaemia in the age group 10-17 years (%)                |             |                |            |             |             |             |                |            |             |             |
| Person  | 909         | 1.3            | 1.4        | 88.9        | 94.8        | 744         | 1.5            | 1.6        | 87.4        | 94.2        |
| Anaemia in the age group 18-59 years (%)                |             |                |            |             |             |             |                |            |             |             |
| Male  | 1420        | 1.2            | 1.3        | 86.5        | 91.7        | 1147        | 1.5            | 1.7        | 86.0        | 92.7        |
| Female  | 1376        | 1.0            | 1.1        | 92.0        | 96.7        | 1060        | 1.4            | 1.4        | 90.8        | 97.1        |
| Anaemia in the age group 60 years and above (%)         |             |                |            |             |             |             |                |            |             |             |
| Male  | 260         | 1.1            | 1.2        | 89.9        | 95.1        | 219         | 1.4            | 1.5        | 88.2        | 94.4        |
| Female  | 241         | 2.4            | 2.6        | 85.4        | 97.0        | 204         | 2.0            | 2.2        | 87.7        | 97.4        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |            |             |             |             |                |            |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |            |             |             |             |                |            |             |             |
| Person  | 3124        | 2.1            | 31.7       | 3.2         | 12.9        | 2490        | 2.7            | 35.6       | 3.3         | 16.0        |
| <b>Hypertension (18 years and above)</b>                |             |                |            |             |             |             |                |            |             |             |
| <b>Above Normal Range</b>                               |             |                |            |             |             |             |                |            |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |            |             |             |             |                |            |             |             |
| Person  | 3132        | 0.4            | 13.8       | 2.2         | 4.0         | 2493        | 0.4            | 17.1       | 1.5         | 3.2         |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |            |             |             |             |                |            |             |             |
| Person  | 3132        | 0.3            | 11.0       | 2.2         | 3.6         | 2493        | 0.4            | 17.4       | 1.5         | 3.2         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |            |             |             |             |                |            |             |             |
| Person  | 3132        | 1.8            | 14.4       | 9.0         | 17.2        | 2493        | 2.2            | 16.1       | 9.7         | 19.8        |
| Iodine content in Household Salt (more than 15 ppm) (%) | <b>1371</b> | <b>5.7</b>     | <b>7.3</b> | <b>62.8</b> | <b>88.0</b> | <b>1100</b> | <b>5.3</b>     | <b>6.7</b> | <b>64.4</b> | <b>87.7</b> |

| Indicators  | Chattarpur  |                |             |             |             |             |                |             |             |             |
|---|-------------|----------------|-------------|-------------|-------------|-------------|----------------|-------------|-------------|-------------|
|   | Total       |                |             |             |             | Rural       |                |             |             |             |
|   | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |             |             |             |             |                |             |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |             |             |             |             |                |             |             |             |
| Person  | 277         | 2.6            | 10.3        | 20.1        | 31.8        | 234         | 2.1            | 8.6         | 19.7        | 28.8        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |             |             |             |             |                |             |             |             |
| Person  | 278         | 4.9            | 8.8         | 44.9        | 66.5        | 235         | 5.6            | 9.7         | 45.1        | 69.7        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |             |             |             |             |                |             |             |             |
| Person  | 299         | 3.1            | 6.6         | 40.0        | 53.6        | 254         | 3.3            | 7.1         | 39.2        | 53.8        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |             |             |             |             |                |             |             |             |
| Person  | 301         | 2.8            | 11.1        | 19.3        | 31.7        | 255         | 2.8            | 12.1        | 17.7        | 30.4        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |             |             |             |             |                |             |             |             |
| Person  | 301         | 3.9            | 17.5        | 14.9        | 32.4        | 255         | 4.3            | 19.3        | 14.4        | 33.7        |
| <b>Nutritional status 5-18 years</b>                    |             |                |             |             |             |             |                |             |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |             |             |             |             |                |             |             |             |
| Person  | 1421        | 4.5            | 15.7        | 19.8        | 39.7        | 1216        | 5.1            | 17.9        | 18.7        | 41.4        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |             |             |             |             |                |             |             |             |
| Person  | 1421        | 0.2            | 22.0        | 0.6         | 1.7         | 1216        | 0.2            | 24.4        | 0.5         | 1.6         |
| <b>Nutritional status 18 years and above</b>            |             |                |             |             |             |             |                |             |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person  | 2664        | 0.6            | 27.7        | 1.3         | 4.1         | 2220        | 0.6            | 25.9        | 1.1         | 4.0         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |             |             |             |             |                |             |             |             |
| Person  | 500         | 0.6            | 22.2        | 1.6         | 4.4         | 417         | 0.7            | 28.2        | 1.3         | 4.7         |
| BMI <18.5 for age 18-59 year (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person  | 2664        | 5.1            | 31.8        | 7.4         | 30.9        | 2220        | 6.1            | 36.5        | 6.8         | 35.0        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |             |             |             |             |                |             |             |             |
| Person  | 500         | 5.7            | 15.6        | 25.0        | 50.2        | 417         | 6.9            | 17.7        | 24.7        | 55.0        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |             |             |             |             |                |             |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |             |             |             |             |                |             |             |             |
| Person  | 219         | 4.3            | 5.9         | 63.1        | 82.3        | 181         | 2.1            | 2.7         | 70.9        | 80.2        |
| Anaemia in the age group 5-9 years (%)                  |             |                |             |             |             |             |                |             |             |             |
| Person  | 374         | 3.2            | 3.7         | 77.3        | 91.7        | 315         | 3.4            | 4.0         | 77.1        | 92.7        |
| Anaemia in the age group 10-17 years (%)                |             |                |             |             |             |             |                |             |             |             |
| Person  | 854         | 2.0            | 2.5         | 77.5        | 86.6        | 733         | 2.3            | 2.7         | 76.6        | 86.6        |
| Anaemia in the age group 18-59 years (%)                |             |                |             |             |             |             |                |             |             |             |
| Male  | 1334        | 1.1            | 1.3         | 87.9        | 93.0        | 1113        | 1.3            | 1.4         | 87.5        | 93.2        |
| Female  | 1175        | 1.7            | 2.0         | 80.3        | 87.9        | 971         | 1.8            | 2.2         | 79.0        | 87.2        |
| Anaemia in the age group 60 years and above (%)         |             |                |             |             |             |             |                |             |             |             |
| Male  | 234         | 2.4            | 2.6         | 84.8        | 95.9        | 195         | 2.9            | 3.1         | 82.5        | 96.0        |
| Female  | 236         | 2.2            | 2.5         | 81.9        | 91.7        | 194         | 2.6            | 3.0         | 80.5        | 92.2        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |             |             |             |             |                |             |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person  | 3107        | 4.4            | 47.0        | 3.1         | 24.7        | -           | -              | -           | -           | -           |
| <b>Hypertension (18 years and above)</b>                |             |                |             |             |             |             |                |             |             |             |
| <b>Above Normal Range</b>                               |             |                |             |             |             |             |                |             |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |             |             |             |             |                |             |             |             |
| Person  | 3174        | 0.5            | 36.9        | 0.6         | 3.1         | 2647        | 0.6            | 45.8        | 0.4         | 3.4         |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |             |             |             |             |                |             |             |             |
| Person  | 3174        | 0.6            | 11.4        | 3.9         | 6.4         | 2647        | 0.6            | 12.6        | 3.5         | 6.2         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |             |             |             |             |                |             |             |             |
| Person  | 3174        | 3.1            | 38.1        | 3.4         | 18.0        | 2647        | 3.6            | 42.0        | 3.3         | 20.8        |
| Iodine content in Household Salt (more than 15 ppm) (%) | <b>1309</b> | <b>10.5</b>    | <b>16.0</b> | <b>40.5</b> | <b>84.6</b> | <b>1120</b> | <b>11.9</b>    | <b>18.6</b> | <b>35.8</b> | <b>84.7</b> |

| Indicators  | Panna       |                |             |             |             |             |                |             |             |             |
|---|-------------|----------------|-------------|-------------|-------------|-------------|----------------|-------------|-------------|-------------|
|   | Total       |                |             |             |             | Rural       |                |             |             |             |
|   | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |             |             |             |             |                |             |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |             |             |             |             |                |             |             |             |
| Person  | 304         | 1.3            | 5.8         | 20.0        | 26.0        | 271         | 1.5            | 6.5         | 19.2        | 25.8        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |             |             |             |             |                |             |             |             |
| Person  | 291         | 3.4            | 7.9         | 35.3        | 50.6        | 258         | 3.2            | 7.8         | 34.2        | 48.7        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |             |             |             |             |                |             |             |             |
| Person  | 330         | 7.8            | 18.9        | 25.2        | 58.9        | 297         | 8.3            | 20.9        | 23.1        | 58.8        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |             |             |             |             |                |             |             |             |
| Person  | 334         | 3.7            | 11.9        | 23.5        | 40.3        | 301         | 4.0            | 12.7        | 23.3        | 41.4        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |             |             |             |             |                |             |             |             |
| Person  | 334         | 2.6            | 14.1        | 13.4        | 25.3        | 301         | 2.7            | 14.0        | 14.1        | 26.6        |
| <b>Nutritional status 5-18 years</b>                    |             |                |             |             |             |             |                |             |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |             |             |             |             |                |             |             |             |
| Person  | 1374        | 1.7            | 8.4         | 16.8        | 24.5        | 1140        | 1.7            | 7.7         | 18.5        | 26.0        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |             |             |             |             |                |             |             |             |
| Person  | 1374        | 0.2            | 40.9        | 0.2         | 1.2         | 1140        | 0.2            | 47.1        | 0.2         | 1.4         |
| <b>Nutritional status 18 years and above</b>            |             |                |             |             |             |             |                |             |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person  | 2458        | 1.4            | 33.9        | 1.9         | 8.7         | 1975        | 1.5            | 43.8        | 1.2         | 9.0         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |             |             |             |             |                |             |             |             |
| Person  | -           | -              | -           | -           | -           | -           | -              | -           | -           | -           |
| BMI <18.5 for age 18-59 year (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person  | 2458        | 0.9            | 8.4         | 9.3         | 13.4        | 1975        | 1.1            | 9.4         | 9.3         | 14.3        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |             |             |             |             |                |             |             |             |
| Person  | 450         | 4.8            | 13.4        | 25.7        | 47.1        | 396         | 5.4            | 14.4        | 26.1        | 50.1        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |             |             |             |             |                |             |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |             |             |             |             |                |             |             |             |
| Person  | 204         | 3.1            | 3.7         | 74.3        | 88.2        | 171         | 3.5            | 4.2         | 72.3        | 87.9        |
| Anaemia in the age group 5-9 years (%)                  |             |                |             |             |             |             |                |             |             |             |
| Person  | 354         | 1.8            | 1.9         | 88.9        | 97.7        | 300         | 2.0            | 2.2         | 87.8        | 97.7        |
| Anaemia in the age group 10-17 years (%)                |             |                |             |             |             |             |                |             |             |             |
| Person  | 824         | 2.3            | 2.6         | 83.0        | 93.4        | 672         | 2.6            | 2.9         | 81.2        | 92.9        |
| Anaemia in the age group 18-59 years (%)                |             |                |             |             |             |             |                |             |             |             |
| Male  | 1220        | 3.6            | 4.4         | 72.7        | 88.5        | 970         | 4.2            | 5.3         | 69.5        | 88.2        |
| Female  | 1108        | 2.5            | 2.8         | 82.1        | 93.7        | 875         | 2.8            | 3.2         | 79.9        | 92.4        |
| Anaemia in the age group 60 years and above (%)         |             |                |             |             |             |             |                |             |             |             |
| Male  | 228         | 2.6            | 2.9         | 82.6        | 94.5        | 198         | 2.9            | 3.2         | 81.9        | 95.3        |
| Female  | 193         | 5.0            | 5.5         | 72.4        | 96.6        | 169         | 5.6            | 6.3         | 69.2        | 96.9        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |             |             |             |             |                |             |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person  | 2885        | 1.2            | 23.0        | 3.2         | 8.7         | 2355        | 1.4            | 29.1        | 2.6         | 9.2         |
| <b>Hypertension (18 years and above)</b>                |             |                |             |             |             |             |                |             |             |             |
| <b>Above Normal Range</b>                               |             |                |             |             |             |             |                |             |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |             |             |             |             |                |             |             |             |
| Person  | 2897        | 0.7            | 14.6        | 3.4         | 6.4         | 2362        | 0.8            | 15.7        | 3.5         | 7.1         |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |             |             |             |             |                |             |             |             |
| Person  | 2897        | 0.4            | 15.0        | 1.7         | 3.3         | 2362        | 0.4            | 18.0        | 1.6         | 3.5         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |             |             |             |             |                |             |             |             |
| Person  | 2897        | 1.3            | 21.2        | 3.7         | 9.4         | 2362        | 0.8            | 14.3        | 4.2         | 7.9         |
| Iodine content in Household Salt (more than 15 ppm) (%) | <b>1255</b> | <b>11.3</b>    | <b>19.0</b> | <b>34.5</b> | <b>80.7</b> | <b>1047</b> | <b>13.7</b>    | <b>23.9</b> | <b>28.1</b> | <b>82.1</b> |

| Indicators  | Sagar       |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 304         | 3.3            | 24.9 | 7.6         | 22.6        | 197         | 3.8            | 22.6 | 10.0        | 26.9        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 253         | 5.1            | 7.2  | 58.3        | 80.3        | 163         | 8.1            | 13.9 | 40.2        | 74.4        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 315         | 7.9            | 15.6 | 33.6        | 66.8        | 202         | 10.0           | 17.4 | 35.6        | 77.1        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 316         | 4.2            | 31.2 | 6.6         | 25.8        | 205         | 5.5            | 31.7 | 8.3         | 32.8        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 316         | 7.2            | 23.7 | 17.2        | 48.3        | 205         | 9.6            | 39.5 | 9.2         | 50.4        |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1714        | 5.9            | 12.1 | 36.2        | 61.4        | 1216        | 3.1            | 5.1  | 53.4        | 66.7        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 1714        | 0.8            | 41.7 | 0.8         | 4.7         | 1216        | 0.4            | 35.7 | 0.5         | 2.2         |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3434        | 2.0            | 39.2 | 2.1         | 12.1        | 2133        | 0.5            | 19.4 | 1.8         | 4.1         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 614         | 2.0            | 31.1 | 3.2         | 12.8        | 424         | 1.0            | 29.8 | 1.7         | 6.5         |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3434        | 4.6            | 12.3 | 27.9        | 48.4        | 2133        | 2.6            | 5.5  | 41.4        | 53.1        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 614         | 4.1            | 7.4  | 46.0        | 64.4        | 424         | 4.6            | 6.8  | 57.2        | 77.8        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 159         | 7.4            | 9.5  | 57.4        | 89.4        | 84          | 1.2            | 1.3  | 87.7        | 92.9        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 421         | 3.1            | 3.5  | 79.9        | 94.3        | 284         | 2.2            | 2.4  | 86.6        | 97.2        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 877         | 2.9            | 3.4  | 77.7        | 90.6        | 624         | 2.5            | 2.8  | 82.1        | 93.1        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1600        | 3.7            | 4.9  | 66.0        | 81.9        | 998         | 5.4            | 7.0  | 63.1        | 86.3        |
| Female  | 1403        | 2.2            | 2.6  | 79.4        | 89.0        | 828         | 2.9            | 3.3  | 80.1        | 92.9        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 252         | 5.2            | 6.5  | 66.8        | 89.2        | 176         | 7.2            | 9.4  | 58.2        | 89.1        |
| Female  | 261         | 3.3            | 4.0  | 74.8        | 89.4        | 176         | 2.5            | 2.8  | 80.9        | 91.8        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3948        | 3.6            | 27.6 | 7.0         | 22.7        | 2508        | 2.9            | 24.5 | 7.0         | 19.9        |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3981        | 1.1            | 21.2 | 3.2         | 8.0         | 2520        | 1.0            | 22.8 | 2.7         | 7.3         |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3981        | 2.1            | 32.1 | 3.3         | 12.9        | 2520        | 1.3            | 28.3 | 2.5         | 8.6         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3981        | 1.0            | 18.9 | 3.5         | 7.9         | 2520        | 1.3            | 23.4 | 3.3         | 9.0         |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1428        | 13.1           | 17.7 | 39.2        | 92.6        | 952         | 18.3           | 24.8 | 26.4        | 95.6        |

| Indicators  | Damoh       |                |             |             |             |             |                |             |             |             |
|---|-------------|----------------|-------------|-------------|-------------|-------------|----------------|-------------|-------------|-------------|
|   | Total       |                |             |             |             | Rural       |                |             |             |             |
|   | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |             |             |             |             |                |             |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |             |             |             |             |                |             |             |             |
| Person  | 306         | 4.1            | 23.1        | 10.3        | 29.0        | 279         | 4.5            | 23.9        | 10.6        | 30.9        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |             |             |             |             |                |             |             |             |
| Person  | 284         | 2.8            | 7.6         | 30.5        | 43.0        | 261         | 3.0            | 8.7         | 28.1        | 41.6        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |             |             |             |             |                |             |             |             |
| Person  | 318         | 4.8            | 16.5        | 19.6        | 41.0        | 290         | 5.0            | 17.7        | 18.6        | 41.1        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |             |             |             |             |                |             |             |             |
| Person  | 320         | 3.7            | 17.0        | 14.4        | 31.0        | 292         | 4.0            | 17.3        | 15.3        | 33.4        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |             |             |             |             |                |             |             |             |
| Person  | 320         | 3.2            | 18.6        | 11.0        | 25.4        | 292         | 3.4            | 21.5        | 9.7         | 25.4        |
| <b>Nutritional status 5-18 years</b>                    |             |                |             |             |             |             |                |             |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |             |             |             |             |                |             |             |             |
| Person  | 1260        | 5.5            | 16.0        | 23.5        | 47.7        | 1080        | 6.3            | 18.5        | 21.6        | 49.0        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |             |             |             |             |                |             |             |             |
| Person  | 1260        | 0.3            | 26.9        | 0.6         | 1.9         | 1080        | 0.3            | 31.9        | 0.4         | 1.9         |
| <b>Nutritional status 18 years and above</b>            |             |                |             |             |             |             |                |             |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person  | 2765        | 1.3            | 22.8        | 3.4         | 9.5         | 2206        | 1.1            | 25.3        | 2.3         | 7.5         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |             |             |             |             |                |             |             |             |
| Person  | 396         | 1.0            | 18.0        | 3.9         | 8.5         | 297         | 1.4            | 28.6        | 2.5         | 9.2         |
| BMI <18.5 for age 18-59 year (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person  | 2765        | 3.0            | 20.5        | 8.9         | 22.7        | 2206        | 3.5            | 21.6        | 9.6         | 25.6        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |             |             |             |             |                |             |             |             |
| Person  | 396         | 4.2            | 11.9        | 26.3        | 45.1        | 297         | 5.5            | 14.6        | 25.9        | 50.3        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |             |             |             |             |                |             |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |             |             |             |             |                |             |             |             |
| Person  | 219         | 5.2            | 7.2         | 58.4        | 81.5        | 195         | 5.7            | 7.8         | 58.4        | 83.9        |
| Anaemia in the age group 5-9 years (%)                  |             |                |             |             |             |             |                |             |             |             |
| Person  | 397         | 2.4            | 2.8         | 78.3        | 89.0        | 356         | 2.2            | 2.6         | 80.2        | 90.3        |
| Anaemia in the age group 10-17 years (%)                |             |                |             |             |             |             |                |             |             |             |
| Person  | 724         | 2.1            | 2.7         | 73.4        | 82.9        | 625         | 2.3            | 2.8         | 75.4        | 85.9        |
| Anaemia in the age group 18-59 years (%)                |             |                |             |             |             |             |                |             |             |             |
| Male  | 1300        | 3.0            | 4.2         | 64.8        | 78.1        | 1051        | 3.6            | 4.9         | 64.1        | 80.0        |
| Female  | 1281        | 2.4            | 2.9         | 76.0        | 86.5        | 1039        | 2.6            | 3.2         | 75.0        | 86.8        |
| Anaemia in the age group 60 years and above (%)         |             |                |             |             |             |             |                |             |             |             |
| Male  | 164         | 3.4            | 4.1         | 74.0        | 89.4        | 119         | 2.3            | 2.5         | 84.9        | 95.4        |
| Female  | 205         | 2.5            | 2.9         | 80.3        | 91.8        | 164         | 2.6            | 3.0         | 81.8        | 93.9        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |             |             |             |             |                |             |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person  | 3121        | 1.1            | 14.0        | 5.9         | 11.0        | 2482        | 1.4            | 17.0        | 5.5         | 11.7        |
| <b>Hypertension (18 years and above)</b>                |             |                |             |             |             |             |                |             |             |             |
| <b>Above Normal Range</b>                               |             |                |             |             |             |             |                |             |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |             |             |             |             |                |             |             |             |
| Person  | 3132        | 0.6            | 13.6        | 3.3         | 6.1         | 2492        | 0.7            | 16.4        | 2.8         | 5.7         |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |             |             |             |             |                |             |             |             |
| Person  | 3132        | 0.8            | 12.1        | 4.9         | 8.5         | 2492        | 1.0            | 14.2        | 4.9         | 9.3         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |             |             |             |             |                |             |             |             |
| Person  | 3132        | 1.0            | 14.2        | 5.3         | 9.9         | 2492        | 1.3            | 17.8        | 4.8         | 10.5        |
| Iodine content in Household Salt (more than 15 ppm) (%) | <b>1360</b> | <b>12.9</b>    | <b>20.4</b> | <b>33.3</b> | <b>85.4</b> | <b>1125</b> | <b>14.9</b>    | <b>24.1</b> | <b>28.4</b> | <b>86.8</b> |

| Indicators  | Satna       |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 403         | 2.2            | 15.4 | 10.1        | 19.7        | 341         | 2.4            | 16.7 | 10.0        | 20.8        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 407         | 3.6            | 13.5 | 19.7        | 35.6        | 344         | 3.4            | 13.3 | 18.7        | 33.6        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 407         | 2.4            | 20.6 | 7.4         | 18.2        | 344         | 1.9            | 18.5 | 6.8         | 15.4        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 407         | 2.2            | 14.9 | 10.6        | 20.4        | 344         | 2.5            | 16.6 | 10.4        | 21.5        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 407         | 0.5            | 29.1 | 0.9         | 3.3         | 344         | 0.6            | 27.3 | 1.2         | 3.9         |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1570        | 3.1            | 10.4 | 23.9        | 37.5        | 1300        | 4.1            | 12.5 | 24.4        | 42.0        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 1570        | 0.3            | 19.1 | 1.1         | 2.5         | 1300        | 0.3            | 25.6 | 0.7         | 2.1         |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2937        | 1.3            | 21.9 | 3.5         | 9.5         | 2317        | 1.0            | 26.9 | 2.2         | 6.9         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 621         | 1.8            | 29.9 | 1.7         | 6.5         | 496         | 2.2            | 50.0 | 1.4         | 13.1        |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2937        | 3.2            | 17.7 | 12.1        | 26.7        | 2317        | 4.2            | 21.4 | 11.8        | 30.8        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 621         | 4.6            | 12.5 | 27.0        | 47.6        | 496         | 5.6            | 14.8 | 26.5        | 51.3        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 273         | 2.3            | 2.4  | 86.9        | 98.5        | 236         | 2.4            | 2.4  | 86.1        | 99.0        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 437         | 1.8            | 1.8  | 89.4        | 99.2        | 375         | 1.1            | 1.1  | 93.0        | 99.6        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 963         | 0.9            | 0.9  | 94.2        | 98.4        | 808         | 1.1            | 1.1  | 93.5        | 99.2        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1350        | 2.3            | 2.4  | 86.4        | 96.9        | 1074        | 2.3            | 2.5  | 86.8        | 97.8        |
| Female  | 1482        | 1.2            | 1.3  | 92.8        | 98.7        | 1181        | 1.5            | 1.5  | 91.3        | 99.2        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 291         | 1.3            | 1.4  | 92.4        | 99.2        | 232         | 1.7            | 1.7  | 90.4        | 99.2        |
| Female  | 297         | 1.0            | 1.0  | 94.0        | 99.0        | 243         | 1.2            | 1.3  | 92.7        | 99.3        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3485        | 0.5            | 9.0  | 4.7         | 7.0         | 2752        | 0.6            | 8.7  | 5.3         | 7.7         |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3463        | 1.2            | 17.2 | 4.7         | 9.9         | 2727        | 1.5            | 25.6 | 3.3         | 9.8         |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3463        | 1.2            | 16.1 | 5.3         | 10.7        | 2727        | 1.3            | 22.8 | 3.6         | 9.5         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3463        | 0.7            | 10.3 | 5.2         | 8.1         | 2727        | 0.8            | 13.7 | 4.2         | 7.6         |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1557        | 4.5            | 5.8  | 66.1        | 85.5        | 1292        | 4.9            | 6.6  | 62.0        | 83.1        |

| Indicators  | Rewa        |                |            |             |             |             |                |             |             |             |
|---|-------------|----------------|------------|-------------|-------------|-------------|----------------|-------------|-------------|-------------|
|   | Total       |                |            |             |             | Rural       |                |             |             |             |
|   | Sample Size | Standard Error | PRSE       | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |            |             |             |             |                |             |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |            |             |             |             |                |             |             |             |
| Person  | 308         | 1.5            | 15.0       | 7.3         | 14.2        | 236         | 1.9            | 19.0        | 6.4         | 14.7        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |            |             |             |             |                |             |             |             |
| Person  | 311         | 3.8            | 9.8        | 31.1        | 47.8        | 240         | 4.7            | 11.7        | 30.0        | 50.2        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |            |             |             |             |                |             |             |             |
| Person  | 320         | 2.7            | 10.7       | 19.5        | 31.3        | 247         | 3.3            | 12.9        | 18.8        | 33.2        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |            |             |             |             |                |             |             |             |
| Person  | 320         | 2.1            | 23.2       | 5.5         | 15.1        | 247         | 2.7            | 28.9        | 4.8         | 17.1        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |            |             |             |             |                |             |             |             |
| Person  | 320         | 1.9            | 31.0       | 3.1         | 12.1        | 247         | 2.3            | 35.0        | 3.0         | 14.0        |
| <b>Nutritional status 5-18 years</b>                    |             |                |            |             |             |             |                |             |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |            |             |             |             |                |             |             |             |
| Person  | 1329        | 3.8            | 13.6       | 20.2        | 36.4        | 1082        | 4.6            | 16.8        | 18.8        | 38.8        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |            |             |             |             |                |             |             |             |
| Person  | 1329        | 0.3            | 18.5       | 1.1         | 2.5         | 1082        | 0.4            | 20.3        | 1.1         | 2.7         |
| <b>Nutritional status 18 years and above</b>            |             |                |            |             |             |             |                |             |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |            |             |             |             |                |             |             |             |
| Person  | 2267        | 1.4            | 17.8       | 5.1         | 11.5        | 1805        | 1.2            | 18.2        | 4.2         | 9.7         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |            |             |             |             |                |             |             |             |
| Person  | 568         | 1.7            | 21.4       | 4.8         | 12.7        | 466         | 2.0            | 23.6        | 4.8         | 14.2        |
| BMI <18.5 for age 18-59 year (%)                        |             |                |            |             |             |             |                |             |             |             |
| Person  | 2267        | 2.1            | 15.3       | 9.9         | 19.5        | 1805        | 2.7            | 19.5        | 8.8         | 21.2        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |            |             |             |             |                |             |             |             |
| Person  | 568         | 3.5            | 13.4       | 19.1        | 34.8        | 466         | 3.9            | 16.0        | 16.9        | 34.4        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |            |             |             |             |                |             |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |            |             |             |             |                |             |             |             |
| Person  | 150         | 4.3            | 4.9        | 74.1        | 93.4        | 117         | 5.2            | 6.1         | 69.8        | 93.2        |
| Anaemia in the age group 5-9 years (%)                  |             |                |            |             |             |             |                |             |             |             |
| Person  | 341         | 3.2            | 3.5        | 81.3        | 96.0        | 274         | 3.4            | 3.8         | 79.3        | 94.8        |
| Anaemia in the age group 10-17 years (%)                |             |                |            |             |             |             |                |             |             |             |
| Person  | 765         | 2.7            | 3.0        | 82.7        | 95.2        | 619         | 3.1            | 3.5         | 79.7        | 94.0        |
| Anaemia in the age group 18-59 years (%)                |             |                |            |             |             |             |                |             |             |             |
| Male  | 1017        | 2.0            | 2.1        | 88.2        | 98.2        | 814         | 2.4            | 2.5         | 86.4        | 97.7        |
| Female  | 1022        | 2.3            | 2.5        | 86.0        | 96.4        | 806         | 2.6            | 2.8         | 83.7        | 95.5        |
| Anaemia in the age group 60 years and above (%)         |             |                |            |             |             |             |                |             |             |             |
| Male  | 288         | 0.9            | 1.0        | 94.5        | 99.0        | 236         | 1.1            | 1.1         | 93.6        | 99.1        |
| Female  | 223         | 1.4            | 1.5        | 90.6        | 97.2        | 187         | 1.6            | 1.7         | 89.4        | 96.4        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |            |             |             |             |                |             |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |            |             |             |             |                |             |             |             |
| Person  | 2733        | 1.0            | 16.9       | 4.3         | 8.9         | 2179        | 1.0            | 18.4        | 3.5         | 7.7         |
| <b>Hypertension (18 years and above)</b>                |             |                |            |             |             |             |                |             |             |             |
| <b>Above Normal Range</b>                               |             |                |            |             |             |             |                |             |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |            |             |             |             |                |             |             |             |
| Person  | 2776        | 0.5            | 25.1       | 1.3         | 3.7         | 2217        | 0.7            | 32.3        | 1.0         | 4.1         |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |            |             |             |             |                |             |             |             |
| Person  | 2776        | 0.6            | 11.1       | 4.3         | 6.9         | 2217        | 0.7            | 13.9        | 3.8         | 6.9         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |            |             |             |             |                |             |             |             |
| Person  | 2776        | 0.8            | 11.4       | 5.5         | 8.9         | 2217        | 1.0            | 13.0        | 5.6         | 9.8         |
| Iodine content in Household Salt (more than 15 ppm) (%) | <b>1182</b> | <b>7.3</b>     | <b>9.2</b> | <b>59.6</b> | <b>90.9</b> | <b>954</b>  | <b>8.2</b>     | <b>10.1</b> | <b>57.5</b> | <b>93.5</b> |

| Indicators  | Umaria      |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 422         | 1.1            | 6.2  | 15.7        | 20.8        | 361         | 1.1            | 6.3  | 15.1        | 20.0        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 374         | 4.7            | 8.1  | 47.0        | 67.8        | 323         | 5.3            | 9.2  | 45.1        | 68.6        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 430         | 3.5            | 7.3  | 39.8        | 55.5        | 367         | 3.8            | 8.1  | 38.5        | 55.5        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 429         | 2.0            | 12.1 | 12.7        | 21.9        | 367         | 2.3            | 14.1 | 11.7        | 22.0        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 429         | 1.9            | 9.6  | 15.9        | 24.6        | 367         | 2.1            | 10.8 | 15.0        | 24.4        |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1549        | 3.0            | 5.4  | 48.7        | 62.0        | 1254        | 3.3            | 5.9  | 49.3        | 64.1        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 1549        | 0.4            | 38.1 | 0.5         | 2.6         | 1254        | 0.3            | 47.6 | 0.3         | 2.1         |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3167        | 1.0            | 33.5 | 1.5         | 6.4         | 2372        | 0.7            | 52.1 | 0.4         | 4.3         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 425         | 1.4            | 43.8 | 1.2         | 8.4         | 329         | 0.3            | 26.7 | 0.8         | 2.2         |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3167        | 3.9            | 12.3 | 23.9        | 41.4        | 2372        | 4.3            | 12.2 | 25.9        | 45.1        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 425         | 3.9            | 10.8 | 27.8        | 45.3        | 329         | 3.5            | 8.6  | 33.0        | 48.7        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 377         | 2.0            | 2.4  | 76.8        | 85.6        | 326         | 2.0            | 2.4  | 75.8        | 84.7        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 492         | 2.3            | 2.5  | 85.8        | 96.6        | 396         | 0.9            | 1.0  | 92.2        | 96.4        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 903         | 1.6            | 1.7  | 88.8        | 96.1        | 750         | 1.1            | 1.2  | 91.9        | 97.2        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1609        | 1.8            | 2.0  | 84.1        | 92.0        | 1209        | 1.6            | 1.7  | 86.0        | 93.0        |
| Female  | 1558        | 1.6            | 1.8  | 88.0        | 95.4        | 1175        | 1.9            | 2.0  | 87.3        | 95.9        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 239         | 1.5            | 1.6  | 88.3        | 95.1        | 186         | 1.4            | 1.5  | 87.9        | 93.9        |
| Female  | 186         | 2.1            | 2.2  | 87.7        | 97.4        | 147         | 2.3            | 2.4  | 86.1        | 96.6        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3491        | 1.1            | 13.8 | 5.9         | 10.9        | 2626        | 1.5            | 19.0 | 5.0         | 11.5        |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3509        | 0.8            | 16.4 | 3.2         | 6.6         | 2640        | 1.0            | 23.6 | 2.5         | 7.0         |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3509        | 0.7            | 12.2 | 4.2         | 7.1         | 2640        | 0.8            | 12.6 | 4.6         | 7.9         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3509        | 1.0            | 20.8 | 3.1         | 7.9         | 2640        | 1.0            | 28.8 | 1.8         | 6.4         |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1452        | 4.4            | 5.4  | 70.0        | 89.8        | 1134        | 4.7            | 6.0  | 66.6        | 87.8        |



| Indicators  | Shahdol     |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 382         | 1.2            | 33.5 | 1.7         | 7.7         | 326         | 1.3            | 35.9 | 1.7         | 8.1         |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 368         | 2.5            | 3.8  | 59.6        | 70.5        | 314         | 3.0            | 4.7  | 56.9        | 70.1        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 395         | 2.2            | 16.3 | 9.1         | 18.8        | 338         | 2.2            | 17.6 | 8.4         | 18.3        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 395         | 1.3            | 36.2 | 1.6         | 7.9         | 338         | 1.5            | 38.9 | 1.6         | 9.1         |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 395         | 3.0            | 9.6  | 25.1        | 38.4        | 338         | 3.1            | 10.2 | 24.0        | 37.7        |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1497        | 2.1            | 7.0  | 25.5        | 34.8        | 1157        | 2.5            | 7.7  | 26.6        | 37.5        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 1497        | 0.4            | 18.9 | 1.5         | 3.5         | 1157        | 0.5            | 20.0 | 1.7         | 4.1         |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3431        | 0.8            | 17.2 | 3.0         | 6.7         | 2367        | 0.7            | 19.0 | 2.5         | 5.7         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 381         | 2.6            | 34.5 | 3.4         | 15.9        | 313         | 3.1            | 42.6 | 2.6         | 18.1        |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3431        | 2.0            | 10.6 | 15.0        | 24.0        | 2367        | 2.8            | 13.9 | 14.4        | 27.1        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 381         | 4.8            | 18.5 | 16.7        | 38.2        | 313         | 5.5            | 19.0 | 18.1        | 42.6        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 221         | 2.4            | 2.9  | 76.0        | 86.7        | 176         | 2.8            | 3.3  | 77.1        | 89.4        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 368         | 1.2            | 1.3  | 91.3        | 96.7        | 286         | 1.0            | 1.0  | 92.9        | 97.6        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 808         | 1.1            | 1.2  | 90.9        | 95.9        | 596         | 0.6            | 0.7  | 93.7        | 96.6        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1517        | 1.2            | 1.3  | 92.6        | 98.6        | 990         | 1.0            | 1.0  | 94.3        | 99.4        |
| Female  | 1456        | 1.5            | 1.6  | 89.6        | 96.5        | 982         | 1.7            | 1.8  | 89.8        | 97.8        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 155         | 0.7            | 0.7  | 93.3        | 99.9        | 117         | 0.9            | 1.0  | 90.9        | 99.9        |
| Female  | 176         | 2.1            | 2.3  | 86.1        | 96.1        | 147         | 2.0            | 2.2  | 87.3        | 97.0        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3673        | 0.7            | 11.9 | 4.6         | 7.8         | 2578        | 0.7            | 11.3 | 4.6         | 7.7         |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3769        | 0.7            | 16.6 | 3.0         | 6.2         | 2654        | 0.7            | 17.2 | 2.8         | 6.1         |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3769        | 0.9            | 10.0 | 7.6         | 11.8        | 2654        | 1.0            | 11.7 | 6.6         | 11.1        |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3769        | 1.1            | 13.9 | 5.6         | 10.3        | 2654        | 1.2            | 14.4 | 6.0         | 11.4        |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1415        | 4.3            | 6.9  | 52.0        | 70.9        | 1067        | 4.8            | 8.4  | 46.1        | 67.2        |

| Indicators  | Sidhi       |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 313         | 3.2            | 21.2 | 9.3         | 24.1        | 259         | 3.5            | 27.1 | 6.8         | 22.8        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 319         | 7.7            | 16.6 | 30.0        | 63.6        | 274         | 8.9            | 18.5 | 29.2        | 67.3        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 323         | 6.6            | 31.0 | 10.0        | 39.7        | 277         | 7.2            | 30.2 | 11.4        | 43.6        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 348         | 2.9            | 13.2 | 16.1        | 29.2        | 274         | 4.7            | 29.5 | 7.9         | 29.6        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 348         | 2.7            | 16.3 | 11.3        | 23.5        | 274         | 3.3            | 18.9 | 11.2        | 26.1        |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1427        | 2.6            | 16.4 | 10.9        | 22.9        | 1212        | 3.0            | 16.1 | 12.7        | 26.3        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 1427        | 0.6            | 32.9 | 0.9         | 3.8         | 1212        | 0.7            | 31.5 | 1.1         | 4.5         |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2461        | 1.5            | 38.4 | 1.5         | 9.1         | 1977        | 0.4            | 24.5 | 0.8         | 2.7         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | -           | -              | -    | -           | -           | -           | -              | -    | -           | -           |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2461        | 1.1            | 17.2 | 4.2         | 9.3         | 1977        | 1.5            | 18.1 | 5.2         | 12.1        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 352         | 2.0            | 7.4  | 22.5        | 31.6        | 305         | 2.3            | 7.3  | 26.8        | 37.2        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 240         | 3.8            | 6.9  | 47.0        | 64.0        | 198         | 4.2            | 7.6  | 46.0        | 65.0        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 404         | 2.8            | 3.8  | 68.0        | 81.0        | 333         | 3.2            | 4.0  | 71.0        | 85.0        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 864         | 2.0            | 2.6  | 72.0        | 81.0        | 746         | 2.4            | 3.1  | 73.0        | 84.0        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1165        | 1.2            | 1.4  | 85.0        | 90.0        | 930         | 1.7            | 1.8  | 85.0        | 93.0        |
| Female  | 1141        | 1.7            | 2.2  | 73.0        | 80.0        | 910         | 2.2            | 2.8  | 74.0        | 84.0        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 175         | 1.6            | 1.8  | 83.0        | 91.0        | 141         | 1.5            | 1.7  | 86.0        | 93.0        |
| Female  | 162         | 2.5            | 2.8  | 81.0        | 93.0        | 153         | 2.6            | 3.0  | 80.0        | 92.0        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2535        | 1.9            | 40.3 | 1.9         | 11.3        | 2008        | 0.9            | 33.6 | 1.2         | 5.4         |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 2802        | 1.9            | 20.8 | 5.5         | 14.1        | 2274        | 1.8            | 25.1 | 4.1         | 12.6        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 2802        | 0.5            | 27.6 | 1.0         | 3.5         | 2274        | 0.6            | 38.5 | 0.7         | 4.0         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 2802        | 1.4            | 17.7 | 5.1         | 11.4        | 2274        | 1.7            | 23.3 | 4.2         | 11.9        |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1345        | 3.9            | 5.4  | 63.1        | 80.7        | 1146        | 4.7            | 6.5  | 61.4        | 82.6        |

| Indicators   | Neemuch     |                |             |             |             |             |                |             |             |             |
|--|-------------|----------------|-------------|-------------|-------------|-------------|----------------|-------------|-------------|-------------|
|  | Total       |                |             |             |             | Rural       |                |             |             |             |
|  | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                        |             |                |             |             |             |             |                |             |             |             |
| Below -2 SD Wasting (Weight for Height) (%)                    |             |                |             |             |             |             |                |             |             |             |
| Person   | 155         | 3.3            | 20.0        | 10.2        | 25.4        | 128         | 4.1            | 24.2        | 9.4         | 28.3        |
| Below -2 SD Stunting (Height for Age) (%)                      |             |                |             |             |             |             |                |             |             |             |
| Person   | 165         | 6.1            | 13.4        | 32.2        | 59.5        | 136         | 6.1            | 12.3        | 36.0        | 63.7        |
| Below -2 SD Underweight (Weight for Age) (%)                   |             |                |             |             |             |             |                |             |             |             |
| Person   | 169         | 5.8            | 16.1        | 24.1        | 50.4        | 138         | 5.4            | 12.9        | 30.4        | 54.9        |
| Below -2 SD Undernourished (BMI for Age) (%)                   |             |                |             |             |             |             |                |             |             |             |
| Person   | 169         | 4.9            | 21.7        | 13.2        | 35.6        | 138         | 6.1            | 26.1        | 12.2        | 39.9        |
| Above 2 SD Overnourished (BMI for Age) (%)                     |             |                |             |             |             |             |                |             |             |             |
| Person   | 169         | 4.1            | 22.4        | 10.6        | 29.6        | 138         | 5.0            | 26.7        | 9.8         | 33.0        |
| <b>Nutritional status 5-18 years</b>                           |             |                |             |             |             |             |                |             |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)                   |             |                |             |             |             |             |                |             |             |             |
| Person   | 832         | 3.3            | 12.1        | 20.4        | 35.3        | 598         | 5.0            | 16.4        | 20.4        | 42.5        |
| Above 2 SD Overnourished (BMI for Age) (%)                     |             |                |             |             |             |             |                |             |             |             |
| Person   | 832         | 1.1            | 28.0        | 2.1         | 7.4         | 598         | 1.4            | 35.5        | 1.8         | 8.7         |
| <b>Nutritional status 18 years and above</b>                   |             |                |             |             |             |             |                |             |             |             |
| BMI ≥25.0 for age 18-59 year (%)                               |             |                |             |             |             |             |                |             |             |             |
| Person   | 2203        | 1.4            | 15.9        | 6.3         | 13.0        | 1452        | 1.5            | 15.9        | 6.5         | 13.3        |
| BMI ≥25.0 for age 60 year and above (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person   | 392         | 1.8            | 23.1        | 4.5         | 12.7        | 288         | 1.5            | 20.4        | 4.6         | 11.6        |
| BMI <18.5 for age 18-59 year (%)                               |             |                |             |             |             |             |                |             |             |             |
| Person   | 2203        | 2.4            | 23.3        | 6.1         | 17.4        | 1452        | 2.0            | 24.3        | 4.6         | 13.8        |
| BMI <18.5 for age 60 year and above (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person   | 392         | 1.0            | 20.6        | 3.0         | 7.6         | 288         | 1.2            | 25.5        | 2.7         | 8.4         |
| <b>Anaemia status by Haemoglobin Level</b>                     |             |                |             |             |             |             |                |             |             |             |
| Anaemia in the age group 6-59 months (%)                       |             |                |             |             |             |             |                |             |             |             |
| Person   | -           | -              | -           | -           | -           | -           | -              | -           | -           | -           |
| Anaemia in the age group 5-9 years (%)                         |             |                |             |             |             |             |                |             |             |             |
| Person   | -           | -              | -           | -           | -           | -           | -              | -           | -           | -           |
| Anaemia in the age group 10-17 years (%)                       |             |                |             |             |             |             |                |             |             |             |
| Person   | 119         | 7.3            | 12.4        | 42.3        | 74.3        | 94          | 7.9            | 13.9        | 38.9        | 73.0        |
| Anaemia in the age group 18-59 years (%)                       |             |                |             |             |             |             |                |             |             |             |
| Male   | 935         | 7.9            | 10.4        | 54.2        | 89.3        | 573         | 7.4            | 10.3        | 52.6        | 85.2        |
| Female   | 804         | 9.7            | 16.9        | 35.7        | 77.2        | 537         | 8.1            | 16.3        | 32.4        | 67.6        |
| Anaemia in the age group 60 years and above (%)                |             |                |             |             |             |             |                |             |             |             |
| Male   | 168         | 6.0            | 7.7         | 62.0        | 89.1        | 104         | 5.2            | 7.1         | 60.3        | 83.6        |
| Female   | 145         | 4.1            | 6.8         | 50.2        | 68.4        | 119         | 3.1            | 5.3         | 51.4        | 65.4        |
| <b>Blood Sugar Level (18 years and above)</b>                  |             |                |             |             |             |             |                |             |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                               |             |                |             |             |             |             |                |             |             |             |
| Person   | 2399        | 1.4            | 11.5        | 9.0         | 15.2        | 1589        | 1.5            | 11.1        | 10.5        | 17.4        |
| <b>Hypertension (18 years and above)</b>                       |             |                |             |             |             |             |                |             |             |             |
| <b>Above Normal Range</b>                                      |             |                |             |             |             |             |                |             |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)              |             |                |             |             |             |             |                |             |             |             |
| Person   | 2496        | 1.5            | 8.9         | 13.4        | 20.1        | 1676        | 1.7            | 9.6         | 14.6        | 22.5        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)              |             |                |             |             |             |             |                |             |             |             |
| Person   | -           | -              | -           | -           | -           | -           | -              | -           | -           | -           |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)              |             |                |             |             |             |             |                |             |             |             |
| Person   | 2496        | 1.0            | 25.1        | 2.3         | 7.1         | 1676        | 1.2            | 27.5        | 2.3         | 8.0         |
| <b>Iodine content in Household Salt (more than 15 ppm) (%)</b> | <b>1155</b> | <b>9.9</b>     | <b>16.4</b> | <b>37.5</b> | <b>79.8</b> | <b>803</b>  | <b>12.2</b>    | <b>21.2</b> | <b>30.5</b> | <b>80.8</b> |

| Indicators  | Mandsaur    |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 299         | 4.4            | 22.9 | 11.1        | 31.6        | 185         | 5.7            | 20.9 | 16.3        | 42.4        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 304         | 3.5            | 5.3  | 56.7        | 72.6        | 186         | 2.4            | 3.8  | 57.5        | 68.5        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 304         | 3.7            | 7.6  | 39.6        | 56.3        | 187         | 5.4            | 10.5 | 38.7        | 62.9        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 322         | 4.4            | 18.3 | 15.5        | 35.8        | 202         | 5.4            | 17.5 | 20.1        | 44.7        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 322         | 1.9            | 25.1 | 4.1         | 13.1        | 202         | 1.9            | 30.0 | 3.1         | 12.4        |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1590        | 1.7            | 5.4  | 28.0        | 35.9        | 1109        | 2.6            | 8.0  | 26.3        | 38.0        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 1590        | 0.2            | 23.6 | 0.6         | 1.7         | 1109        | 0.2            | 30.5 | 0.4         | 1.5         |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3596        | 0.8            | 15.9 | 3.5         | 7.2         | 2533        | 0.5            | 15.4 | 2.3         | 4.5         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 518         | 2.0            | 15.1 | 9.2         | 18.1        | 382         | 1.6            | 16.9 | 6.4         | 13.8        |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3596        | 1.8            | 7.4  | 20.1        | 28.0        | 2533        | 2.1            | 9.4  | 18.0        | 27.5        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 518         | 3.9            | 19.2 | 12.9        | 30.7        | 382         | 4.4            | 19.3 | 14.5        | 34.5        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 60          | 7.2            | 10.6 | 49.9        | 82.3        | 39          | 9.1            | 15.7 | 36.9        | 76.5        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 189         | 6.3            | 8.1  | 59.8        | 88.4        | 153         | 7.5            | 10.3 | 52.9        | 86.8        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 560         | 5.5            | 7.0  | 63.5        | 89.0        | 446         | 6.9            | 9.1  | 56.9        | 88.2        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1118        | 3.2            | 3.8  | 76.3        | 91.4        | 888         | 4.2            | 4.9  | 73.0        | 93.0        |
| Female  | 1192        | 4.5            | 5.7  | 66.2        | 86.3        | 923         | 5.8            | 7.6  | 60.9        | 87.0        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 174         | 2.8            | 3.1  | 82.3        | 96.0        | 146         | 3.4            | 3.7  | 79.9        | 97.0        |
| Female  | 157         | 7.5            | 9.7  | 56.6        | 90.0        | 119         | 10.2           | 13.5 | 46.9        | 91.6        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3587        | 1.0            | 12.1 | 6.5         | 11.2        | 2556        | 1.5            | 19.7 | 4.9         | 12.1        |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 4330        | 1.2            | 8.2  | 12.5        | 18.1        | 2992        | 1.8            | 12.8 | 10.7        | 19.0        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 4330        | 1.6            | 26.9 | 3.1         | 10.6        | 2992        | 1.1            | 25.6 | 2.5         | 7.9         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 4330        | 0.6            | 13.2 | 3.3         | 6.0         | 2992        | 0.6            | 12.4 | 3.4         | 5.9         |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1519        | 10.5           | 14.5 | 44.5        | 89.9        | 1075        | 15.0           | 22.3 | 30.6        | 90.5        |

| Indicators  | Ratlam      |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 174         | 9.3            | 25.7 | 18.5        | 58.3        | 118         | 12.5           | 32.3 | 16.0        | 67.3        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 179         | 5.3            | 15.8 | 22.7        | 46.1        | 121         | 6.9            | 16.8 | 26.7        | 57.1        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 192         | 3.4            | 8.2  | 34.1        | 49.4        | 133         | 4.4            | 8.9  | 39.5        | 59.1        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 194         | 9.8            | 25.5 | 19.7        | 61.6        | 135         | 13.0           | 32.0 | 16.8        | 69.7        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 194         | 4.3            | 44.6 | 3.4         | 24.8        | -           | -              | -    | -           | -           |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1153        | 3.6            | 10.5 | 26.4        | 42.5        | 796         | 4.7            | 14.1 | 23.8        | 44.8        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 1153        | 0.6            | 44.6 | 0.5         | 3.6         | -           | -              | -    | -           | -           |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2740        | 1.3            | 16.4 | 5.5         | 11.7        | 1662        | 1.5            | 24.8 | 3.5         | 10.7        |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 482         | 3.5            | 26.7 | 7.1         | 23.5        | 281         | 3.1            | 36.7 | 3.6         | 18.7        |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2740        | 3.3            | 11.7 | 21.4        | 36.3        | 1662        | 5.0            | 19.3 | 16.4        | 38.9        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 482         | 3.6            | 15.3 | 16.3        | 32.5        | 281         | 5.5            | 25.0 | 12.1        | 37.0        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 36          | 4.5            | 5.2  | 72.8        | 93.9        | -           | -              | -    | -           | -           |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 108         | 7.1            | 8.4  | 61.7        | 95.0        | 73          | 7.9            | 9.1  | 58.3        | 96.6        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 372         | 2.7            | 3.1  | 80.7        | 93.1        | 238         | 1.5            | 1.6  | 87.6        | 94.4        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1168        | 2.4            | 2.6  | 85.3        | 96.8        | 731         | 0.7            | 0.8  | 93.6        | 97.0        |
| Female  | 1095        | 3.1            | 3.5  | 79.9        | 94.3        | 621         | 1.8            | 1.9  | 86.2        | 94.3        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 213         | 2.6            | 2.9  | 83.1        | 95.3        | 117         | 3.4            | 3.7  | 80.1        | 98.1        |
| Female  | 195         | 4.3            | 4.9  | 74.1        | 94.2        | 114         | 1.9            | 2.0  | 87.3        | 96.1        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2938        | 1.2            | 10.3 | 9.2         | 14.6        | 1777        | 1.2            | 11.4 | 8.1         | 13.5        |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3386        | 1.4            | 9.7  | 11.9        | 18.4        | 2011        | 2.2            | 16.1 | 9.3         | 19.2        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3386        | 1.8            | 29.5 | 3.0         | 11.4        | 2011        | 2.9            | 43.1 | 2.5         | 16.9        |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3386        | 1.9            | 28.8 | 3.4         | 12.3        | 2011        | 2.0            | 31.2 | 3.1         | 12.7        |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1269        | 9.9            | 13.0 | 48.2        | 91.8        | 789         | 14.8           | 21.5 | 31.7        | 91.2        |

| Indicators  | Ujjain      |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 324         | 3.2            | 12.3 | 19.3        | 33.6        | 202         | 4.2            | 16.6 | 17.1        | 36.2        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 306         | 3.8            | 7.2  | 43.5        | 60.3        | 191         | 2.1            | 3.4  | 56.0        | 65.3        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 321         | 3.9            | 8.8  | 35.3        | 52.6        | 200         | 4.4            | 9.4  | 37.4        | 57.2        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 350         | 3.6            | 12.3 | 22.2        | 38.6        | 215         | 5.2            | 18.4 | 18.1        | 41.4        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 350         | 1.4            | 23.5 | 3.6         | 10.3        | 215         | 2.2            | 38.0 | 2.4         | 13.1        |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1288        | 6.2            | 15.0 | 28.3        | 55.6        | 799         | 9.9            | 24.0 | 21.9        | 64.1        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 1288        | 0.4            | 39.7 | 0.4         | 2.6         | -           | -              | -    | -           | -           |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3199        | 0.9            | 15.6 | 4.0         | 8.2         | 1775        | 0.2            | 5.0  | 3.6         | 4.5         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 627         | 2.7            | 20.0 | 8.4         | 20.8        | 361         | 0.5            | 7.7  | 5.2         | 7.4         |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3199        | 2.2            | 6.1  | 31.5        | 41.5        | 1775        | 2.6            | 7.4  | 29.6        | 41.5        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 627         | 2.9            | 11.2 | 20.0        | 33.1        | 361         | 2.1            | 7.9  | 21.9        | 31.3        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 117         | 5.4            | 6.6  | 65.7        | 90.0        | 73          | 5.6            | 6.7  | 67.4        | 93.6        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 219         | 4.4            | 5.2  | 72.5        | 93.0        | 141         | 2.6            | 2.9  | 82.2        | 94.5        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 543         | 4.9            | 5.8  | 70.1        | 93.0        | 344         | 5.7            | 6.5  | 68.3        | 96.6        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1172        | 4.7            | 5.6  | 70.2        | 91.6        | 696         | 6.1            | 7.3  | 65.1        | 93.3        |
| Female  | 1210        | 3.9            | 4.4  | 76.2        | 94.4        | 656         | 6.1            | 6.9  | 66.6        | 96.1        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 264         | 3.6            | 4.2  | 76.2        | 92.9        | 173         | 5.0            | 5.9  | 69.3        | 92.2        |
| Female  | 216         | 3.3            | 3.7  | 78.6        | 94.0        | 108         | 5.0            | 5.9  | 69.7        | 92.9        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3324        | 1.3            | 11.8 | 8.5         | 14.5        | 1969        | 1.4            | 18.1 | 5.0         | 11.4        |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3914        | 1.3            | 9.4  | 11.4        | 17.4        | 2195        | 1.9            | 15.8 | 8.5         | 17.2        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3914        | 0.8            | 14.9 | 3.7         | 7.3         | 2195        | 1.1            | 24.2 | 2.6         | 7.9         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3914        | 0.8            | 13.0 | 4.8         | 8.6         | 2195        | 1.3            | 20.2 | 4.1         | 10.2        |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1270        | 15.4           | 19.2 | 31.4        | 97.2        | 773         | 25.0           | 38.4 | 13.4        | 95.7        |

| Indicators  | Shajapur    |                |             |             |             |             |                |             |             |             |
|---|-------------|----------------|-------------|-------------|-------------|-------------|----------------|-------------|-------------|-------------|
|   | Total       |                |             |             |             | Rural       |                |             |             |             |
|   | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |             |             |             |             |                |             |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |             |             |             |             |                |             |             |             |
| Person  | 308         | 4.4            | 28.6        | 7.8         | 28.1        | 231         | 5.2            | 29.5        | 8.8         | 32.7        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |             |             |             |             |                |             |             |             |
| Person  | 321         | 5.0            | 9.2         | 43.2        | 65.6        | 239         | 6.4            | 10.7        | 45.0        | 73.1        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |             |             |             |             |                |             |             |             |
| Person  | 323         | 4.2            | 12.1        | 26.2        | 45.2        | 241         | 5.9            | 16.8        | 23.1        | 49.2        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |             |             |             |             |                |             |             |             |
| Person  | 332         | 4.7            | 21.6        | 12.9        | 34.1        | 250         | 5.6            | 23.6        | 13.5        | 38.7        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |             |             |             |             |                |             |             |             |
| Person  | 332         | 2.1            | 25.6        | 4.6         | 14.7        | 250         | 3.2            | 33.8        | 4.3         | 19.6        |
| <b>Nutritional status 5-18 years</b>                    |             |                |             |             |             |             |                |             |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |             |             |             |             |                |             |             |             |
| Person  | 1327        | 3.0            | 8.5         | 29.2        | 42.8        | 1044        | 3.3            | 9.4         | 28.1        | 42.9        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |             |             |             |             |                |             |             |             |
| Person  | -           | -              | -           | -           | -           | -           | -              | -           | -           | -           |
| <b>Nutritional status 18 years and above</b>            |             |                |             |             |             |             |                |             |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person  | 3073        | 1.5            | 17.6        | 5.6         | 12.5        | 2445        | 1.3            | 15.7        | 5.7         | 11.6        |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |             |             |             |             |                |             |             |             |
| Person  | 622         | 1.8            | 18.5        | 6.5         | 14.9        | 545         | 2.1            | 23.5        | 5.2         | 15.1        |
| BMI <18.5 for age 18-59 year (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person  | 3073        | 2.8            | 21.3        | 8.0         | 20.9        | 2445        | 3.7            | 28.3        | 6.8         | 24.2        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |             |             |             |             |                |             |             |             |
| Person  | 622         | 2.9            | 29.2        | 5.0         | 18.5        | 545         | 3.4            | 29.1        | 5.9         | 21.8        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |             |             |             |             |                |             |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |             |             |             |             |                |             |             |             |
| Person  | 143         | 8.5            | 14.3        | 39.6        | 76.4        | 102         | 10.7           | 17.4        | 36.4        | 81.4        |
| Anaemia in the age group 5-9 years (%)                  |             |                |             |             |             |             |                |             |             |             |
| Person  | 288         | 8.9            | 14.2        | 41.6        | 80.0        | 218         | 9.1            | 14.3        | 41.7        | 80.8        |
| Anaemia in the age group 10-17 years (%)                |             |                |             |             |             |             |                |             |             |             |
| Person  | 674         | 6.0            | 10.8        | 41.8        | 68.3        | 517         | 6.6            | 12.6        | 37.6        | 66.6        |
| Anaemia in the age group 18-59 years (%)                |             |                |             |             |             |             |                |             |             |             |
| Male  | 1477        | 6.4            | 9.6         | 51.4        | 79.9        | 1194        | 7.2            | 11.3        | 46.6        | 78.1        |
| Female  | 1280        | 7.1            | 11.6        | 44.8        | 76.1        | 989         | 6.9            | 12.1        | 41.2        | 71.3        |
| Anaemia in the age group 60 years and above (%)         |             |                |             |             |             |             |                |             |             |             |
| Male  | 284         | 7.5            | 10.3        | 53.7        | 87.0        | 250         | 8.9            | 12.5        | 48.0        | 86.8        |
| Female  | 233         | 6.3            | 9.3         | 52.4        | 80.3        | 200         | 7.0            | 10.4        | 49.8        | 80.4        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |             |             |             |             |                |             |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person  | 3365        | 2.0            | 18.2        | 7.1         | 16.2        | 2711        | 1.4            | 15.8        | 6.2         | 12.7        |
| <b>Hypertension (18 years and above)</b>                |             |                |             |             |             |             |                |             |             |             |
| <b>Above Normal Range</b>                               |             |                |             |             |             |             |                |             |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |             |             |             |             |                |             |             |             |
| Person  | 3482        | 0.7            | 4.2         | 15.3        | 18.5        | 2809        | 0.8            | 4.3         | 16.1        | 19.5        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |             |             |             |             |                |             |             |             |
| Person  | 3482        | 0.8            | 19.1        | 2.7         | 6.5         | 2809        | 1.0            | 20.9        | 2.9         | 7.4         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |             |             |             |             |                |             |             |             |
| Person  | 3482        | 1.2            | 25.5        | 2.7         | 8.6         | 2809        | 1.1            | 30.4        | 1.9         | 7.3         |
| Iodine content in Household Salt (more than 15 ppm) (%) | <b>1389</b> | <b>10.1</b>    | <b>18.6</b> | <b>32.3</b> | <b>75.1</b> | <b>1119</b> | <b>10.1</b>    | <b>21.4</b> | <b>26.2</b> | <b>68.8</b> |

| Indicators  | Dewas       |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 395         | 3.5            | 12.8 | 19.9        | 35.6        | 283         | 4.6            | 16.8 | 18.1        | 38.6        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 321         | 3.8            | 8.0  | 39.3        | 56.4        | 241         | 4.9            | 10.2 | 37.5        | 59.4        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 410         | 4.8            | 8.7  | 44.2        | 65.5        | 295         | 6.7            | 12.8 | 37.6        | 67.0        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 411         | 3.3            | 14.8 | 15.6        | 30.5        | 295         | 4.3            | 19.4 | 13.9        | 33.2        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 411         | 1.8            | 39.7 | 1.8         | 10.6        | 295         | 2.3            | 48.2 | 1.6         | 13.9        |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1784        | 2.3            | 5.0  | 40.8        | 51.3        | 1294        | 3.3            | 7.1  | 39.1        | 53.9        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 1784        | 0.6            | 16.2 | 2.5         | 5.3         | 1294        | 0.6            | 20.4 | 2.0         | 4.9         |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3371        | 0.6            | 29.8 | 1.1         | 4.1         | 2309        | 0.9            | 34.5 | 1.2         | 5.7         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 491         | 1.2            | 25.8 | 2.6         | 8.5         | 346         | 1.8            | 29.6 | 3.1         | 11.7        |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3371        | 2.6            | 7.1  | 30.7        | 42.3        | 2309        | 2.9            | 7.7  | 31.0        | 43.9        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 491         | 7.0            | 14.6 | 32.7        | 63.2        | 346         | 8.6            | 19.1 | 27.0        | 64.0        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 106         | 8.6            | 15.3 | 36.5        | 73.5        | 82          | 11.2           | 19.8 | 31.6        | 78.4        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 286         | 7.8            | 11.7 | 47.6        | 81.8        | 210         | 8.8            | 12.3 | 48.6        | 87.0        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 608         | 3.4            | 5.4  | 55.5        | 70.9        | 418         | 4.8            | 7.9  | 49.6        | 71.1        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1230        | 2.9            | 3.8  | 67.7        | 80.7        | 821         | 4.1            | 5.7  | 62.1        | 80.6        |
| Female  | 1252        | 3.6            | 5.6  | 55.1        | 71.1        | 860         | 3.5            | 5.5  | 55.8        | 71.5        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 160         | 5.8            | 9.1  | 50.3        | 76.1        | 117         | 8.2            | 13.4 | 41.6        | 77.1        |
| Female  | 141         | 7.8            | 11.1 | 50.2        | 84.4        | 105         | 11.0           | 16.0 | 41.0        | 87.6        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3255        | 3.7            | 48.9 | 2.4         | 21.2        | -           | -              | -    | -           | -           |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3938        | 2.9            | 18.1 | 10.6        | 23.9        | 2686        | 4.3            | 27.8 | 8.0         | 27.8        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3938        | 1.1            | 24.6 | 2.5         | 7.4         | 2686        | 1.2            | 29.1 | 2.1         | 7.8         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3938        | 1.2            | 22.2 | 3.2         | 8.7         | 2686        | 1.7            | 30.6 | 2.8         | 11.1        |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1201        | 10.9           | 17.4 | 36.9        | 82.9        | 842         | 16.3           | 32.5 | 18.7        | 81.6        |



| Indicators  | Jhabua      |                |             |             |             |             |                |             |             |             |
|---|-------------|----------------|-------------|-------------|-------------|-------------|----------------|-------------|-------------|-------------|
|   | Total       |                |             |             |             | Rural       |                |             |             |             |
|   | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |             |             |             |             |                |             |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |             |             |             |             |                |             |             |             |
| Person  | 192         | 2.2            | 18.7        | 7.8         | 18.1        | 170         | 2.3            | 18.8        | 7.9         | 18.5        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |             |             |             |             |                |             |             |             |
| Person  | 199         | 6.4            | 12.1        | 38.5        | 66.8        | 176         | 7.0            | 13.4        | 36.7        | 67.3        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |             |             |             |             |                |             |             |             |
| Person  | 235         | 4.1            | 13.2        | 22.8        | 41.3        | 212         | 4.4            | 14.2        | 22.0        | 41.8        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |             |             |             |             |                |             |             |             |
| Person  | 250         | 1.3            | 17.1        | 5.3         | 11.5        | 227         | 1.4            | 16.3        | 5.8         | 12.1        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |             |             |             |             |                |             |             |             |
| Person  | 250         | 3.0            | 12.3        | 18.4        | 32.1        | 227         | 3.1            | 13.0        | 17.8        | 32.0        |
| <b>Nutritional status 5-18 years</b>                    |             |                |             |             |             |             |                |             |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |             |             |             |             |                |             |             |             |
| Person  | 1753        | 1.3            | 5.5         | 20.7        | 26.6        | 1595        | 1.4            | 5.9         | 20.4        | 26.6        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |             |             |             |             |                |             |             |             |
| Person  | 1753        | 0.2            | 26.6        | 0.3         | 1.2         | 1595        | 0.2            | 26.5        | 0.4         | 1.2         |
| <b>Nutritional status 18 years and above</b>            |             |                |             |             |             |             |                |             |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person  | 2669        | 0.2            | 20.6        | 0.6         | 1.5         | 2352        | 0.2            | 25.2        | 0.4         | 1.3         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |             |             |             |             |                |             |             |             |
| Person  | -           | -              | -           | -           | -           | -           | -              | -           | -           | -           |
| BMI <18.5 for age 18-59 year (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person  | 2669        | 2.1            | 6.1         | 30.2        | 39.8        | 2352        | 2.3            | 6.6         | 29.9        | 40.4        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |             |             |             |             |                |             |             |             |
| Person  | 298         | 5.5            | 13.1        | 30.6        | 55.2        | 246         | 5.9            | 14.8        | 27.7        | 54.0        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |             |             |             |             |                |             |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |             |             |             |             |                |             |             |             |
| Person  | 139         | 6.8            | 10.3        | 49.4        | 79.3        | 125         | 7.2            | 11.2        | 47.0        | 78.6        |
| Anaemia in the age group 5-9 years (%)                  |             |                |             |             |             |             |                |             |             |             |
| Person  | 530         | 4.5            | 6.1         | 62.4        | 82.4        | 491         | 4.6            | 6.4         | 60.7        | 81.5        |
| Anaemia in the age group 10-17 years (%)                |             |                |             |             |             |             |                |             |             |             |
| Person  | 979         | 4.1            | 5.5         | 64.3        | 82.9        | 885         | 4.3            | 5.8         | 62.8        | 81.9        |
| Anaemia in the age group 18-59 years (%)                |             |                |             |             |             |             |                |             |             |             |
| Male  | 1309        | 5.2            | 6.2         | 67.8        | 91.4        | 1159        | 5.5            | 6.8         | 65.5        | 90.6        |
| Female  | 1132        | 5.4            | 7.2         | 61.4        | 85.7        | 988         | 5.8            | 7.8         | 58.8        | 84.5        |
| Anaemia in the age group 60 years and above (%)         |             |                |             |             |             |             |                |             |             |             |
| Male  | 141         | 4.4            | 5.1         | 73.3        | 94.1        | 119         | 4.9            | 5.8         | 70.7        | 93.8        |
| Female  | 131         | 3.1            | 3.9         | 71.6        | 85.6        | 106         | 3.4            | 4.4         | 68.8        | 84.1        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |             |             |             |             |                |             |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person  | 2731        | 0.8            | 8.2         | 8.2         | 11.9        | 2378        | 0.9            | 9.1         | 7.8         | 11.7        |
| <b>Hypertension (18 years and above)</b>                |             |                |             |             |             |             |                |             |             |             |
| <b>Above Normal Range</b>                               |             |                |             |             |             |             |                |             |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |             |             |             |             |                |             |             |             |
| Person  | 2837        | 2.9            | 16.8        | 11.7        | 25.0        | 2479        | 3.2            | 17.5        | 12.1        | 26.6        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |             |             |             |             |                |             |             |             |
| Person  | 2837        | 0.1            | 33.4        | 0.2         | 0.8         | 2479        | 0.1            | 35.2        | 0.2         | 0.9         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |             |             |             |             |                |             |             |             |
| Person  | 2837        | 1.3            | 15.4        | 6.0         | 12.1        | 2479        | 1.5            | 17.0        | 5.8         | 12.5        |
| Iodine content in Household Salt (more than 15 ppm) (%) | <b>1144</b> | <b>12.9</b>    | <b>28.3</b> | <b>20.5</b> | <b>73.1</b> | <b>999</b>  | <b>13.1</b>    | <b>30.5</b> | <b>18.4</b> | <b>71.7</b> |

| Indicators  | Dhar        |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 117         | 5.2            | 22.0 | 14.1        | 37.6        | 88          | 4.9            | 20.6 | 14.5        | 36.6        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 117         | 6.6            | 12.7 | 37.0        | 65.8        | 91          | 8.1            | 15.2 | 35.2        | 70.2        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 148         | 3.7            | 8.6  | 34.5        | 50.8        | 115         | 4.3            | 10.5 | 31.6        | 50.7        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 179         | 3.2            | 25.1 | 7.1         | 21.8        | 144         | 3.5            | 26.7 | 6.9         | 23.1        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 179         | 8.3            | 38.4 | 8.4         | 45.7        | 144         | 10.2           | 40.3 | 9.1         | 53.5        |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1444        | 2.0            | 5.6  | 32.1        | 41.3        | 1222        | 2.3            | 6.1  | 32.3        | 42.6        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 1444        | 1.1            | 34.5 | 1.4         | 6.7         | 1222        | 1.3            | 35.3 | 1.6         | 7.9         |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3012        | 0.4            | 22.4 | 1.0         | 2.8         | 2414        | 0.4            | 27.7 | 0.8         | 2.8         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 391         | 0.9            | 28.1 | 1.6         | 5.7         | 335         | 0.8            | 46.4 | 0.6         | 5.1         |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3012        | 2.3            | 5.6  | 35.4        | 45.6        | 2414        | 2.7            | 6.7  | 35.0        | 47.4        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 391         | 3.7            | 9.0  | 33.3        | 49.9        | 335         | 4.0            | 9.2  | 34.9        | 52.9        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 54          | 3.9            | 6.4  | 51.9        | 69.6        | 36          | 5.5            | 10.0 | 43.0        | 67.5        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 222         | 4.9            | 6.9  | 59.5        | 81.6        | 166         | 6.3            | 9.0  | 54.3        | 82.5        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 557         | 3.0            | 3.8  | 72.4        | 86.0        | 460         | 3.6            | 4.6  | 68.8        | 84.9        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1202        | 2.1            | 2.4  | 81.2        | 90.8        | 946         | 2.5            | 2.9  | 80.1        | 91.6        |
| Female  | 1039        | 2.6            | 3.2  | 73.5        | 85.3        | 809         | 3.2            | 4.1  | 70.8        | 85.5        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 167         | 4.6            | 5.1  | 73.9        | 97.2        | 148         | 5.2            | 5.7  | 70.9        | 97.3        |
| Female  | 117         | 3.6            | 4.5  | 69.8        | 86.0        | 94          | 4.0            | 5.0  | 68.8        | 86.8        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2958        | 0.9            | 11.4 | 6.3         | 10.6        | 2387        | 0.8            | 12.5 | 4.9         | 8.6         |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3490        | 1.3            | 8.7  | 11.9        | 17.6        | 2816        | 1.5            | 11.2 | 10.1        | 16.8        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3490        | 1.7            | 28.2 | 3.2         | 11.3        | 2816        | 1.7            | 32.0 | 2.5         | 10.6        |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3490        | 0.8            | 15.1 | 3.6         | 7.2         | 2816        | 0.9            | 19.8 | 2.9         | 7.0         |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1193        | 12.0           | 23.4 | 26.2        | 75.8        | 975         | 12.6           | 26.9 | 21.9        | 73.4        |

| Indicators  | Indore      |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 507         | 1.3            | 18.6 | 4.4         | 10.2        | 139         | 0.9            | 7.1  | 10.3        | 14.2        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 508         | 6.4            | 10.6 | 45.6        | 73.9        | 137         | 2.8            | 5.7  | 43.2        | 56.0        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 530         | 5.5            | 15.0 | 25.3        | 49.7        | 146         | 3.0            | 8.8  | 28.0        | 41.7        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 566         | 1.2            | 20.7 | 3.5         | 9.0         | 171         | 0.9            | 10.4 | 7.1         | 11.3        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 566         | 2.4            | 22.7 | 6.3         | 17.5        | 171         | 1.1            | 9.7  | 9.2         | 14.2        |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 2023        | 0.7            | 2.3  | 26.6        | 29.6        | 634         | 0.6            | 1.8  | 32.7        | 35.4        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 2023        | 0.9            | 43.3 | 0.8         | 5.5         | 634         | 0.0            | 4.2  | 0.5         | 0.6         |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 5235        | 1.1            | 8.1  | 11.2        | 16.1        | 1407        | 1.4            | 16.9 | 5.7         | 12.2        |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 711         | 1.7            | 7.5  | 18.9        | 26.5        | 183         | 1.0            | 8.2  | 9.7         | 14.0        |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 5235        | 2.8            | 8.3  | 27.4        | 39.9        | 1407        | 1.0            | 2.3  | 41.8        | 46.3        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 711         | 1.8            | 7.4  | 20.7        | 29.0        | 183         | 3.3            | 11.7 | 21.3        | 36.1        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 151         | 4.8            | 6.0  | 66.9        | 88.5        | 43          | 1.1            | 1.9  | 57.7        | 62.9        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 264         | 4.8            | 6.5  | 61.7        | 83.2        | 98          | 4.5            | 9.5  | 37.0        | 57.0        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 575         | 4.7            | 6.7  | 58.4        | 79.2        | 186         | 1.0            | 2.4  | 39.2        | 43.6        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1052        | 5.2            | 7.3  | 58.4        | 81.5        | 298         | 2.2            | 4.9  | 40.7        | 50.8        |
| Female  | 1568        | 6.3            | 8.3  | 59.1        | 87.4        | 421         | 2.3            | 4.7  | 43.8        | 54.3        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 214         | 5.8            | 7.4  | 62.5        | 88.4        | 64          | 3.4            | 8.1  | 34.5        | 49.9        |
| Female  | 184         | 7.5            | 10.4 | 52.8        | 85.9        | 46          | 3.0            | 13.5 | 16.2        | 29.8        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 5271        | 0.7            | 6.5  | 8.8         | 11.7        | 1514        | 0.0            | 0.7  | 5.5         | 5.7         |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 6088        | 0.6            | 4.2  | 13.8        | 16.7        | 1646        | 0.2            | 0.9  | 17.0        | 17.7        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 6088        | 1.9            | 27.8 | 3.6         | 12.8        | 1646        | 2.0            | 21.4 | 5.7         | 14.9        |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 6088        | 0.1            | 18.4 | 0.4         | 0.9         | 1646        | 0.0            | 3.6  | 0.5         | 0.6         |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1704        | 1.4            | 1.6  | 85.6        | 92.1        | 493         | 1.5            | 2.2  | 65.4        | 72.2        |

| Indicators  | West Nimar  |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 598         | 2.4            | 16.2 | 10.2        | 21.2        | 516         | 2.7            | 18.4 | 9.6         | 21.9        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 637         | 5.3            | 7.4  | 58.5        | 82.2        | 547         | 6.0            | 8.1  | 58.4        | 85.2        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 633         | 3.5            | 5.4  | 57.4        | 73.3        | 543         | 3.8            | 5.7  | 57.9        | 75.1        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 642         | 3.3            | 17.3 | 12.7        | 27.7        | 552         | 3.9            | 20.5 | 11.7        | 29.3        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 642         | 1.4            | 21.9 | 3.9         | 10.5        | 552         | 1.5            | 25.2 | 3.3         | 10.1        |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1379        | 2.4            | 10.7 | 17.4        | 28.1        | 1165        | 2.7            | 11.7 | 17.8        | 30.1        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 1379        | 0.6            | 30.5 | 1.0         | 4.1         | 1165        | 0.7            | 30.2 | 1.2         | 4.8         |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3049        | 0.6            | 15.7 | 2.8         | 5.8         | 2492        | 0.7            | 19.3 | 2.3         | 5.4         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 380         | 1.4            | 28.9 | 2.4         | 9.0         | 312         | 1.1            | 39.8 | 1.2         | 7.0         |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3049        | 1.0            | 6.0  | 14.8        | 19.4        | 2492        | 1.2            | 6.8  | 14.4        | 19.6        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 380         | 2.4            | 16.5 | 9.9         | 20.9        | 312         | 2.7            | 20.3 | 8.2         | 20.4        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 37          | 13.1           | 20.9 | 32.0        | 85.5        | -           | -              | -    | -           | -           |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 192         | 9.0            | 11.9 | 50.9        | 90.8        | 145         | 12.4           | 17.4 | 38.6        | 90.5        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 583         | 11.0           | 16.1 | 40.5        | 87.0        | 481         | 13.1           | 20.1 | 33.5        | 87.1        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1279        | 10.6           | 13.5 | 46.9        | 93.6        | 1042        | 12.8           | 17.2 | 38.9        | 93.3        |
| Female  | 1256        | 10.2           | 13.9 | 45.7        | 89.8        | 1010        | 12.4           | 18.0 | 37.5        | 89.3        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 172         | 9.9            | 11.9 | 50.1        | 95.7        | 140         | 11.6           | 14.4 | 43.8        | 95.5        |
| Female  | 146         | 9.9            | 15.0 | 41.7        | 84.2        | 115         | 12.3           | 19.5 | 34.0        | 84.8        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3096        | 1.3            | 13.7 | 7.0         | 13.0        | 2519        | 1.7            | 19.6 | 5.4         | 13.1        |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3304        | 2.5            | 13.8 | 12.9        | 24.0        | 2713        | 3.0            | 17.8 | 11.1        | 24.8        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3304        | 0.7            | 13.9 | 3.6         | 6.8         | 2713        | 0.7            | 16.9 | 2.9         | 6.2         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3304        | 1.2            | 20.4 | 3.8         | 9.5         | 2713        | 1.0            | 17.5 | 3.9         | 8.6         |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1409        | 15.1           | 20.3 | 32.6        | 94.6        | 1190        | 17.4           | 24.2 | 26.7        | 94.7        |

| Indicators   | Barwani     |                |             |             |             |             |                |             |             |             |
|--|-------------|----------------|-------------|-------------|-------------|-------------|----------------|-------------|-------------|-------------|
|  | Total       |                |             |             |             | Rural       |                |             |             |             |
|  | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                        |             |                |             |             |             |             |                |             |             |             |
| Below -2 SD Wasting (Weight for Height) (%)                    |             |                |             |             |             |             |                |             |             |             |
| Person   | 381         | 7.9            | 28.4        | 13.6        | 48.3        | 364         | 7.9            | 29.5        | 12.9        | 47.9        |
| Below -2 SD Stunting (Height for Age) (%)                      |             |                |             |             |             |             |                |             |             |             |
| Person   | 437         | 6.3            | 11.1        | 42.7        | 70.6        | 413         | 6.8            | 11.6        | 42.9        | 72.9        |
| Below -2 SD Underweight (Weight for Age) (%)                   |             |                |             |             |             |             |                |             |             |             |
| Person   | 436         | 5.6            | 8.5         | 52.2        | 76.9        | 412         | 5.8            | 8.9         | 51.4        | 77.3        |
| Below -2 SD Undernourished (BMI for Age) (%)                   |             |                |             |             |             |             |                |             |             |             |
| Person   | 463         | 8.0            | 21.7        | 21.2        | 56.2        | 433         | 7.9            | 22.5        | 19.8        | 54.4        |
| Above 2 SD Overnourished (BMI for Age) (%)                     |             |                |             |             |             |             |                |             |             |             |
| Person   | 463         | 1.0            | 23.0        | 2.5         | 7.2         | 433         | 1.0            | 24.5        | 2.3         | 6.8         |
| <b>Nutritional status 5-18 years</b>                           |             |                |             |             |             |             |                |             |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)                   |             |                |             |             |             |             |                |             |             |             |
| Person   | 1689        | 4.6            | 10.4        | 34.4        | 55.0        | 1517        | 5.0            | 11.2        | 33.7        | 55.9        |
| Above 2 SD Overnourished (BMI for Age) (%)                     |             |                |             |             |             |             |                |             |             |             |
| Person   | 1689        | 1.2            | 30.7        | 2.0         | 7.9         | 1517        | 1.3            | 33.7        | 1.8         | 8.0         |
| <b>Nutritional status 18 years and above</b>                   |             |                |             |             |             |             |                |             |             |             |
| BMI ≥25.0 for age 18-59 year (%)                               |             |                |             |             |             |             |                |             |             |             |
| Person   | 2821        | 0.6            | 18.3        | 2.1         | 4.9         | 2384        | 0.6            | 20.4        | 1.9         | 4.8         |
| BMI ≥25.0 for age 60 year and above (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person   | 389         | 3.0            | 24.3        | 7.0         | 20.8        | 298         | 3.5            | 30.1        | 5.7         | 22.1        |
| BMI <18.5 for age 18-59 year (%)                               |             |                |             |             |             |             |                |             |             |             |
| Person   | 2821        | 3.1            | 10.3        | 23.5        | 37.5        | 2384        | 3.7            | 12.2        | 22.5        | 39.0        |
| BMI <18.5 for age 60 year and above (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person   | 389         | 2.9            | 11.7        | 18.7        | 31.7        | 298         | 3.0            | 11.2        | 20.3        | 33.6        |
| <b>Anaemia status by Haemoglobin Level</b>                     |             |                |             |             |             |             |                |             |             |             |
| Anaemia in the age group 6-59 months (%)                       |             |                |             |             |             |             |                |             |             |             |
| Person   | -           | -              | -           | -           | -           | -           | -              | -           | -           | -           |
| Anaemia in the age group 5-9 years (%)                         |             |                |             |             |             |             |                |             |             |             |
| Person   | 136         | 7.8            | 13.0        | 41.8        | 75.8        | 115         | 9.2            | 15.4        | 38.5        | 77.8        |
| Anaemia in the age group 10-17 years (%)                       |             |                |             |             |             |             |                |             |             |             |
| Person   | 456         | 10.9           | 17.1        | 37.7        | 83.6        | 410         | 11.7           | 18.4        | 35.7        | 84.6        |
| Anaemia in the age group 18-59 years (%)                       |             |                |             |             |             |             |                |             |             |             |
| Male   | 999         | 5.3            | 7.3         | 59.5        | 83.4        | 842         | 5.3            | 7.2         | 59.7        | 83.4        |
| Female   | 939         | 6.0            | 9.1         | 51.1        | 77.7        | 780         | 6.7            | 10.0        | 50.5        | 80.2        |
| Anaemia in the age group 60 years and above (%)                |             |                |             |             |             |             |                |             |             |             |
| Male   | 149         | 4.4            | 5.6         | 66.4        | 86.3        | 107         | 4.1            | 5.2         | 69.0        | 87.7        |
| Female   | 118         | 6.5            | 10.9        | 44.6        | 73.3        | 90          | 8.1            | 12.6        | 45.0        | 80.3        |
| <b>Blood Sugar Level (18 years and above)</b>                  |             |                |             |             |             |             |                |             |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                               |             |                |             |             |             |             |                |             |             |             |
| Person   | 2775        | 1.0            | 12.0        | 6.1         | 10.5        | 2306        | 1.1            | 13.7        | 5.8         | 10.8        |
| <b>Hypertension (18 years and above)</b>                       |             |                |             |             |             |             |                |             |             |             |
| <b>Above Normal Range</b>                                      |             |                |             |             |             |             |                |             |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)              |             |                |             |             |             |             |                |             |             |             |
| Person   | 3219        | 1.8            | 11.6        | 11.9        | 20.1        | 2679        | 2.1            | 14.4        | 10.3        | 19.8        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)              |             |                |             |             |             |             |                |             |             |             |
| Person   | 3219        | 1.6            | 26.7        | 3.2         | 10.5        | 2679        | 1.6            | 30.7        | 2.6         | 10.4        |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)              |             |                |             |             |             |             |                |             |             |             |
| Person   | 3219        | 1.6            | 29.8        | 2.7         | 10.5        | 2679        | 1.4            | 33.5        | 1.9         | 8.7         |
| <b>Iodine content in Household Salt (more than 15 ppm) (%)</b> | <b>1117</b> | <b>15.0</b>    | <b>23.0</b> | <b>29.6</b> | <b>89.4</b> | <b>930</b>  | <b>16.9</b>    | <b>28.8</b> | <b>22.7</b> | <b>87.5</b> |

| Indicators  | East Nimar  |                |            |             |             |             |                |             |             |             |
|---|-------------|----------------|------------|-------------|-------------|-------------|----------------|-------------|-------------|-------------|
|   | Total       |                |            |             |             | Rural       |                |             |             |             |
|   | Sample Size | Standard Error | PRSE       | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |            |             |             |             |                |             |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |            |             |             |             |                |             |             |             |
| Person  | 544         | 3.6            | 39.4       | 3.7         | 21.3        | 415         | 4.6            | 38.7        | 4.7         | 26.5        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |            |             |             |             |                |             |             |             |
| Person  | 594         | 6.4            | 15.2       | 28.6        | 56.7        | 459         | 8.5            | 20.9        | 23.7        | 60.6        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |            |             |             |             |                |             |             |             |
| Person  | 614         | 5.6            | 18.7       | 19.0        | 44.0        | 471         | 6.3            | 23.9        | 14.8        | 43.0        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |            |             |             |             |                |             |             |             |
| Person  | 625         | 7.8            | 37.4       | 8.2         | 43.2        | 477         | 9.9            | 40.2        | 8.9         | 52.1        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |            |             |             |             |                |             |             |             |
| Person  | 625         | 1.7            | 29.0       | 3.0         | 11.0        | 477         | 1.4            | 20.4        | 4.2         | 10.5        |
| <b>Nutritional status 5-18 years</b>                    |             |                |            |             |             |             |                |             |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |            |             |             |             |                |             |             |             |
| Person  | 1518        | 2.4            | 7.0        | 28.8        | 39.7        | 1274        | 2.5            | 6.9         | 31.2        | 42.7        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |            |             |             |             |                |             |             |             |
| Person  | 1518        | 0.2            | 39.8       | 0.2         | 1.0         | -           | -              | -           | -           | -           |
| <b>Nutritional status 18 years and above</b>            |             |                |            |             |             |             |                |             |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |            |             |             |             |                |             |             |             |
| Person  | 3178        | 1.7            | 30.4       | 2.8         | 10.9        | 2600        | 2.1            | 39.2        | 2.1         | 12.4        |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |            |             |             |             |                |             |             |             |
| Person  | 539         | 1.8            | 18.3       | 6.3         | 14.4        | 449         | 1.8            | 23.1        | 4.7         | 13.2        |
| BMI <18.5 for age 18-59 year (%)                        |             |                |            |             |             |             |                |             |             |             |
| Person  | 3178        | 3.6            | 14.5       | 17.4        | 33.5        | 2600        | 4.3            | 18.4        | 15.1        | 34.6        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |            |             |             |             |                |             |             |             |
| Person  | 539         | 4.1            | 24.3       | 9.5         | 28.2        | 449         | 4.9            | 28.3        | 8.9         | 31.4        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |            |             |             |             |                |             |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |            |             |             |             |                |             |             |             |
| Person  | 267         | 7.9            | 11.5       | 49.2        | 83.9        | 251         | 8.4            | 12.2        | 47.6        | 84.2        |
| Anaemia in the age group 5-9 years (%)                  |             |                |            |             |             |             |                |             |             |             |
| Person  | 350         | 5.6            | 7.3        | 62.2        | 87.6        | 319         | 6.0            | 7.8         | 60.3        | 87.2        |
| Anaemia in the age group 10-17 years (%)                |             |                |            |             |             |             |                |             |             |             |
| Person  | 695         | 5.6            | 7.6        | 59.1        | 84.0        | 593         | 6.6            | 9.2         | 54.6        | 83.8        |
| Anaemia in the age group 18-59 years (%)                |             |                |            |             |             |             |                |             |             |             |
| Male  | 1382        | 6.7            | 8.7        | 58.7        | 88.8        | 1145        | 8.4            | 11.1        | 52.4        | 89.5        |
| Female  | 1311        | 4.8            | 6.3        | 63.8        | 85.4        | 1068        | 6.0            | 8.0         | 59.2        | 85.9        |
| Anaemia in the age group 60 years and above (%)         |             |                |            |             |             |             |                |             |             |             |
| Male  | 254         | 5.6            | 6.9        | 65.6        | 91.3        | 213         | 6.8            | 8.7         | 59.1        | 89.8        |
| Female  | 210         | 6.9            | 8.7        | 59.8        | 91.3        | 178         | 8.3            | 10.5        | 54.8        | 92.0        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |            |             |             |             |                |             |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |            |             |             |             |                |             |             |             |
| Person  | 3401        | 3.5            | 33.3       | 4.8         | 21.3        | 2823        | 4.1            | 39.2        | 4.2         | 24.1        |
| <b>Hypertension (18 years and above)</b>                |             |                |            |             |             |             |                |             |             |             |
| <b>Above Normal Range</b>                               |             |                |            |             |             |             |                |             |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |            |             |             |             |                |             |             |             |
| Person  | 3466        | 3.2            | 20.3       | 9.8         | 24.3        | 2875        | 3.9            | 25.2        | 8.6         | 26.6        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |            |             |             |             |                |             |             |             |
| Person  | 3466        | 0.5            | 13.2       | 2.8         | 5.1         | 2875        | 0.6            | 14.6        | 2.8         | 5.4         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |            |             |             |             |                |             |             |             |
| Person  | 3466        | 1.8            | 27.8       | 3.4         | 11.7        | 2875        | 2.0            | 33.7        | 2.8         | 12.7        |
| Iodine content in Household Salt (more than 15 ppm) (%) | <b>1328</b> | <b>7.1</b>     | <b>8.9</b> | <b>59.1</b> | <b>90.9</b> | <b>1087</b> | <b>8.8</b>     | <b>11.6</b> | <b>51.3</b> | <b>90.2</b> |

| Indicators  | Rajgarh     |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 269         | 1.9            | 9.3  | 16.6        | 25.3        | 213         | 2.0            | 8.7  | 18.8        | 27.8        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 273         | 5.8            | 11.9 | 36.1        | 61.7        | 217         | 5.7            | 11.4 | 37.4        | 62.5        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 273         | 8.8            | 20.0 | 25.9        | 63.7        | 217         | 8.8            | 18.9 | 27.9        | 65.6        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 274         | 1.9            | 7.9  | 19.7        | 28.2        | 218         | 2.2            | 8.4  | 21.5        | 31.5        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | -           | -              | -    | -           | -           | -           | -              | -    | -           | -           |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1119        | 3.6            | 16.2 | 14.9        | 31.1        | 972         | 4.0            | 17.8 | 14.7        | 32.9        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | -           | -              | -    | -           | -           | -           | -              | -    | -           | -           |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2552        | 0.4            | 20.2 | 1.1         | 2.7         | 2120        | 0.4            | 21.8 | 1.1         | 3.1         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 488         | 1.4            | 27.9 | 2.6         | 9.1         | 419         | 1.5            | 32.5 | 2.2         | 9.5         |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2552        | 2.8            | 31.1 | 4.3         | 17.5        | 2120        | 3.2            | 33.7 | 4.3         | 19.7        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 488         | 3.2            | 32.2 | 4.6         | 19.6        | 419         | 3.5            | 33.0 | 4.9         | 21.4        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 159         | 6.4            | 18.6 | 21.8        | 50.2        | 123         | 7.6            | 22.4 | 19.2        | 52.4        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 288         | 4.1            | 7.9  | 42.3        | 60.4        | 251         | 4.4            | 8.6  | 40.9        | 60.5        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 653         | 4.0            | 7.9  | 41.9        | 60.0        | 573         | 4.4            | 8.9  | 40.0        | 59.7        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1265        | 4.0            | 6.1  | 57.2        | 75.3        | 1061        | 4.7            | 7.0  | 55.8        | 76.8        |
| Female  | 1133        | 5.0            | 9.9  | 39.3        | 61.6        | 933         | 5.8            | 11.5 | 37.6        | 63.2        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 254         | 5.4            | 8.0  | 54.2        | 78.3        | 218         | 6.1            | 9.1  | 51.6        | 78.5        |
| Female  | 219         | 5.7            | 12.3 | 33.9        | 59.2        | 189         | 6.4            | 13.7 | 32.9        | 61.0        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2923        | 2.0            | 17.5 | 7.7         | 16.9        | 2424        | 1.6            | 17.4 | 6.1         | 13.4        |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3015        | 1.9            | 13.4 | 10.5        | 19.2        | 2515        | 1.6            | 11.9 | 10.1        | 17.3        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3015        | 0.5            | 13.6 | 2.9         | 5.3         | 2515        | 0.6            | 17.4 | 2.3         | 5.1         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3015        | 1.1            | 20.3 | 3.4         | 8.4         | 2515        | 0.9            | 19.5 | 3.1         | 7.4         |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1200        | 17.2           | 27.1 | 24.5        | 90.4        | 1017        | 19.4           | 31.9 | 19.8        | 90.7        |

| Indicators  | Vidisha     |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 382         | 4.1            | 14.3 | 20.3        | 38.8        | 271         | 4.1            | 15.3 | 18.8        | 37.4        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 343         | 6.4            | 11.8 | 39.6        | 67.7        | 232         | 6.2            | 11.3 | 40.7        | 67.9        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 397         | 3.0            | 7.0  | 36.9        | 50.6        | 272         | 3.7            | 9.1  | 32.4        | 48.9        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 412         | 4.7            | 17.2 | 17.9        | 38.9        | 283         | 4.0            | 16.3 | 16.8        | 35.0        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 412         | 3.2            | 15.1 | 14.7        | 29.1        | 283         | 3.8            | 17.0 | 15.0        | 32.4        |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1659        | 2.6            | 9.0  | 23.7        | 35.6        | 964         | 2.8            | 8.5  | 26.7        | 39.2        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 1659        | 0.9            | 17.0 | 3.5         | 7.5         | 964         | 0.9            | 19.1 | 3.0         | 7.1         |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3235        | 2.5            | 22.3 | 6.8         | 18.3        | 1758        | 3.1            | 41.9 | 2.9         | 18.2        |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 499         | 2.9            | 21.3 | 8.2         | 21.5        | 288         | 3.4            | 34.7 | 4.4         | 20.6        |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3235        | 3.4            | 17.9 | 12.5        | 27.9        | 1758        | 3.9            | 17.1 | 15.5        | 33.0        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 499         | 3.6            | 16.8 | 14.2        | 30.5        | 288         | 2.9            | 10.7 | 21.1        | 34.1        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 268         | 5.9            | 7.5  | 62.1        | 88.6        | 175         | 6.4            | 7.8  | 62.8        | 92.0        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 438         | 2.3            | 2.7  | 81.4        | 92.1        | 271         | 2.7            | 3.1  | 80.0        | 92.6        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 910         | 3.9            | 5.0  | 68.4        | 86.1        | 550         | 5.0            | 6.7  | 62.4        | 85.0        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1530        | 5.6            | 8.3  | 54.1        | 79.2        | 858         | 8.3            | 12.7 | 45.4        | 81.5        |
| Female  | 1364        | 3.1            | 3.7  | 75.6        | 89.6        | 753         | 4.6            | 5.6  | 68.7        | 89.4        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 235         | 4.6            | 5.5  | 69.8        | 90.7        | 141         | 5.9            | 7.6  | 62.1        | 88.7        |
| Female  | 225         | 4.3            | 5.0  | 72.8        | 92.7        | 134         | 5.5            | 6.5  | 68.0        | 93.4        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | -           | -              | -    | -           | -           | -           | -              | -    | -           | -           |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3667        | 0.6            | 13.2 | 3.1         | 5.7         | 2015        | 0.9            | 20.4 | 2.7         | 6.8         |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3667        | 1.3            | 18.3 | 4.7         | 10.7        | 2015        | 0.8            | 16.1 | 3.5         | 7.2         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3667        | 2.1            | 37.0 | 2.5         | 13.0        | 2015        | 3.2            | 47.5 | 2.2         | 18.6        |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1297        | 10.4           | 16.4 | 38.5        | 82.6        | 691         | 4.5            | 4.7  | 70.1        | 99.3        |



| Indicators   | Bhopal      |                |            |             |             |             |                |            |             |             |
|--|-------------|----------------|------------|-------------|-------------|-------------|----------------|------------|-------------|-------------|
|  | Total       |                |            |             |             | Rural       |                |            |             |             |
|  | Sample Size | Standard Error | PRSE       | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE       | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                        |             |                |            |             |             |             |                |            |             |             |
| Below -2 SD Wasting (Weight for Height) (%)                    |             |                |            |             |             |             |                |            |             |             |
| Person   | 542         | 2.1            | 19.5       | 6.8         | 16.4        | 85          | 0.2            | 1.0        | 20.5        | 21.5        |
| Below -2 SD Stunting (Height for Age) (%)                      |             |                |            |             |             |             |                |            |             |             |
| Person   | 548         | 7.3            | 13.2       | 38.9        | 71.0        | 96          | 0.2            | 0.7        | 29.3        | 30.3        |
| Below -2 SD Underweight (Weight for Age) (%)                   |             |                |            |             |             |             |                |            |             |             |
| Person   | 557         | 4.3            | 9.3        | 36.4        | 55.6        | 96          | 0.2            | 0.6        | 41.3        | 42.4        |
| Below -2 SD Undernourished (BMI for Age) (%)                   |             |                |            |             |             |             |                |            |             |             |
| Person   | 569         | 2.7            | 18.1       | 9.6         | 21.7        | 98          | 0.2            | 0.7        | 31.1        | 32.1        |
| Above 2 SD Overnourished (BMI for Age) (%)                     |             |                |            |             |             |             |                |            |             |             |
| Person   | 569         | 0.9            | 12.9       | 4.9         | 8.8         | 98          | 0.1            | 1.6        | 7.6         | 8.2         |
| <b>Nutritional status 5-18 years</b>                           |             |                |            |             |             |             |                |            |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)                   |             |                |            |             |             |             |                |            |             |             |
| Person   | 1376        | 2.2            | 7.2        | 25.9        | 35.8        | 309         | 0.1            | 0.4        | 36.5        | 37.1        |
| Above 2 SD Overnourished (BMI for Age) (%)                     |             |                |            |             |             |             |                |            |             |             |
| Person   | 1376        | 0.4            | 18.6       | 1.3         | 2.9         | 309         | 0.0            | 1.7        | 2.3         | 2.5         |
| <b>Nutritional status 18 years and above</b>                   |             |                |            |             |             |             |                |            |             |             |
| BMI ≥25.0 for age 18-59 year (%)                               |             |                |            |             |             |             |                |            |             |             |
| Person   | 3886        | 1.0            | 7.8        | 10.4        | 14.7        | 663         | 0.0            | 0.9        | 3.8         | 3.9         |
| BMI ≥25.0 for age 60 year and above (%)                        |             |                |            |             |             |             |                |            |             |             |
| Person   | 546         | 1.7            | 8.5        | 16.0        | 23.4        | 126         | 0.1            | 1.5        | 6.8         | 7.3         |
| BMI <18.5 for age 18-59 year (%)                               |             |                |            |             |             |             |                |            |             |             |
| Person   | 3886        | 1.6            | 5.7        | 25.2        | 32.5        | 663         | 0.1            | 0.2        | 39.9        | 40.4        |
| BMI <18.5 for age 60 year and above (%)                        |             |                |            |             |             |             |                |            |             |             |
| Person   | 546         | 1.4            | 6.5        | 18.0        | 24.2        | 126         | 0.2            | 0.6        | 32.5        | 33.4        |
| <b>Anaemia status by Haemoglobin Level</b>                     |             |                |            |             |             |             |                |            |             |             |
| Anaemia in the age group 6-59 months (%)                       |             |                |            |             |             |             |                |            |             |             |
| Person   | 57          | 6.8            | 9.1        | 56.7        | 87.0        | -           | -              | -          | -           | -           |
| Anaemia in the age group 5-9 years (%)                         |             |                |            |             |             |             |                |            |             |             |
| Person   | 179         | 4.1            | 4.8        | 72.7        | 91.4        | 35          | 0.2            | 0.2        | 94.1        | 94.9        |
| Anaemia in the age group 10-17 years (%)                       |             |                |            |             |             |             |                |            |             |             |
| Person   | 486         | 3.7            | 4.6        | 71.1        | 87.8        | 108         | 0.1            | 0.1        | 92.0        | 92.5        |
| Anaemia in the age group 18-59 years (%)                       |             |                |            |             |             |             |                |            |             |             |
| Male   | 1248        | 4.9            | 6.2        | 65.8        | 87.8        | 253         | 0.1            | 0.1        | 89.7        | 90.1        |
| Female   | 1394        | 2.8            | 3.4        | 74.9        | 87.6        | 225         | 0.1            | 0.1        | 87.5        | 87.9        |
| Anaemia in the age group 60 years and above (%)                |             |                |            |             |             |             |                |            |             |             |
| Male   | 200         | 2.8            | 3.3        | 78.4        | 91.4        | 52          | 0.1            | 0.1        | 95.7        | 96.3        |
| Female   | 207         | 5.5            | 6.9        | 64.0        | 88.7        | 49          | 0.2            | 0.2        | 87.8        | 88.8        |
| <b>Blood Sugar Level (18 years and above)</b>                  |             |                |            |             |             |             |                |            |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                               |             |                |            |             |             |             |                |            |             |             |
| Person   | 3836        | 1.0            | 9.7        | 8.1         | 12.6        | 701         | 0.1            | 0.4        | 15.0        | 15.3        |
| <b>Hypertension (18 years and above)</b>                       |             |                |            |             |             |             |                |            |             |             |
| <b>Above Normal Range</b>                                      |             |                |            |             |             |             |                |            |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)              |             |                |            |             |             |             |                |            |             |             |
| Person   | 4701        | 2.3            | 13.9       | 12.1        | 22.7        | 832         | 0.1            | 0.4        | 13.8        | 14.1        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)              |             |                |            |             |             |             |                |            |             |             |
| Person   | 4701        | 0.3            | 5.5        | 4.8         | 6.2         | 832         | 0.0            | 0.7        | 5.9         | 6.0         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)              |             |                |            |             |             |             |                |            |             |             |
| Person   | 4701        | 0.4            | 8.9        | 3.6         | 5.4         | 832         | 0.0            | 0.5        | 8.8         | 9.1         |
| <b>Iodine content in Household Salt (more than 15 ppm) (%)</b> | <b>1481</b> | <b>0.3</b>     | <b>0.3</b> | <b>97.9</b> | <b>99.2</b> | <b>278</b>  | <b>0.1</b>     | <b>0.1</b> | <b>93.4</b> | <b>93.8</b> |

| Indicators  | Sehore      |                |            |             |             |             |                |             |             |             |
|---|-------------|----------------|------------|-------------|-------------|-------------|----------------|-------------|-------------|-------------|
|   | Total       |                |            |             |             | Rural       |                |             |             |             |
|   | Sample Size | Standard Error | PRSE       | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |            |             |             |             |                |             |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |            |             |             |             |                |             |             |             |
| Person  | 421         | 3.1            | 23.7       | 7.5         | 21.7        | 345         | 3.7            | 28.3        | 6.7         | 23.7        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |            |             |             |             |                |             |             |             |
| Person  | 451         | 5.9            | 11.9       | 36.6        | 62.8        | 373         | 6.7            | 12.7        | 37.7        | 67.2        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |            |             |             |             |                |             |             |             |
| Person  | 449         | 4.0            | 8.7        | 37.1        | 54.9        | 369         | 4.0            | 8.3         | 39.1        | 56.8        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |            |             |             |             |                |             |             |             |
| Person  | 462         | 2.9            | 13.6       | 15.4        | 28.5        | 381         | 3.3            | 15.4        | 14.8        | 29.6        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |            |             |             |             |                |             |             |             |
| Person  | 462         | 0.9            | 24.2       | 2.1         | 6.2         | 381         | 1.0            | 25.3        | 2.2         | 6.8         |
| <b>Nutritional status 5-18 years</b>                    |             |                |            |             |             |             |                |             |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |            |             |             |             |                |             |             |             |
| Person  | 1181        | 3.0            | 10.6       | 22.1        | 35.6        | 869         | 4.1            | 13.7        | 21.7        | 40.2        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |            |             |             |             |                |             |             |             |
| Person  | -           | -              | -          | -           | -           | -           | -              | -           | -           | -           |
| <b>Nutritional status 18 years and above</b>            |             |                |            |             |             |             |                |             |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |            |             |             |             |                |             |             |             |
| Person  | 2441        | 1.0            | 12.8       | 5.9         | 10.5        | 1883        | 1.2            | 16.0        | 5.3         | 10.9        |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |            |             |             |             |                |             |             |             |
| Person  | 354         | 2.7            | 27.5       | 5.3         | 18.2        | 308         | 3.1            | 32.9        | 4.4         | 19.4        |
| BMI <18.5 for age 18-59 year (%)                        |             |                |            |             |             |             |                |             |             |             |
| Person  | 2441        | 3.7            | 17.8       | 13.6        | 30.3        | 1883        | 4.6            | 22.2        | 12.1        | 32.8        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |            |             |             |             |                |             |             |             |
| Person  | 354         | 3.9            | 20.1       | 12.1        | 30.0        | 308         | 4.2            | 23.1        | 10.5        | 29.8        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |            |             |             |             |                |             |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |            |             |             |             |                |             |             |             |
| Person  | 124         | 7.0            | 11.2       | 45.5        | 76.1        | 80          | 7.9            | 12.4        | 44.8        | 79.0        |
| Anaemia in the age group 5-9 years (%)                  |             |                |            |             |             |             |                |             |             |             |
| Person  | 224         | 3.7            | 4.8        | 67.5        | 83.9        | 157         | 3.2            | 4.1         | 71.2        | 85.9        |
| Anaemia in the age group 10-17 years (%)                |             |                |            |             |             |             |                |             |             |             |
| Person  | 496         | 4.4            | 6.0        | 61.7        | 81.5        | 294         | 4.6            | 6.1         | 63.1        | 83.7        |
| Anaemia in the age group 18-59 years (%)                |             |                |            |             |             |             |                |             |             |             |
| Male  | 722         | 2.8            | 3.2        | 79.2        | 91.8        | 533         | 2.4            | 2.8         | 81.9        | 93.1        |
| Female  | 857         | 3.0            | 3.9        | 69.8        | 83.4        | 625         | 2.5            | 3.2         | 72.1        | 83.5        |
| Anaemia in the age group 60 years and above (%)         |             |                |            |             |             |             |                |             |             |             |
| Male  | 134         | 3.1            | 3.5        | 79.5        | 93.7        | 115         | 3.1            | 3.6         | 78.5        | 92.9        |
| Female  | 118         | 5.5            | 6.9        | 64.7        | 89.8        | 102         | 5.3            | 6.6         | 65.9        | 90.2        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |            |             |             |             |                |             |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |            |             |             |             |                |             |             |             |
| Person  | 2470        | 1.4            | 11.5       | 9.6         | 16.2        | 1925        | 0.7            | 5.5         | 11.2        | 14.4        |
| <b>Hypertension (18 years and above)</b>                |             |                |            |             |             |             |                |             |             |             |
| <b>Above Normal Range</b>                               |             |                |            |             |             |             |                |             |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |            |             |             |             |                |             |             |             |
| Person  | 2906        | 3.1            | 17.3       | 11.9        | 26.0        | 2272        | 3.5            | 20.2        | 10.7        | 26.6        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |            |             |             |             |                |             |             |             |
| Person  | 2906        | 0.7            | 23.8       | 1.6         | 4.8         | 2272        | 0.7            | 26.5        | 1.3         | 4.5         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |            |             |             |             |                |             |             |             |
| Person  | 2906        | 1.9            | 27.0       | 3.8         | 12.7        | 2272        | 2.2            | 26.3        | 4.5         | 14.6        |
| Iodine content in Household Salt (more than 15 ppm) (%) | <b>1065</b> | <b>6.8</b>     | <b>7.9</b> | <b>63.1</b> | <b>95.9</b> | <b>832</b>  | <b>8.8</b>     | <b>10.3</b> | <b>54.6</b> | <b>96.3</b> |

| Indicators  | Raisen      |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 322         | 6.2            | 26.7 | 12.4        | 39.1        | 259         | 7.3            | 29.5 | 12.3        | 43.5        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 269         | 6.3            | 9.9  | 48.9        | 75.7        | 214         | 7.2            | 11.9 | 44.3        | 74.7        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 340         | 4.1            | 6.8  | 50.7        | 68.3        | 274         | 4.7            | 7.5  | 51.7        | 71.9        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 345         | 5.9            | 28.3 | 10.8        | 36.4        | 279         | 7.0            | 31.6 | 10.5        | 40.6        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 345         | 3.7            | 13.9 | 19.3        | 35.3        | 279         | 4.3            | 16.5 | 17.7        | 36.1        |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1132        | 2.9            | 8.5  | 28.3        | 41.0        | 882         | 3.1            | 9.4  | 26.5        | 39.9        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 1132        | 1.1            | 45.5 | 0.9         | 6.7         | 882         | 1.4            | 46.3 | 1.1         | 8.0         |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2198        | 2.6            | 21.6 | 7.2         | 19.2        | 1571        | 2.6            | 23.6 | 6.3         | 18.4        |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 344         | 3.8            | 37.5 | 4.3         | 22.5        | 267         | 2.7            | 34.5 | 3.5         | 16.5        |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2198        | 2.7            | 11.7 | 17.8        | 29.9        | 1571        | 2.9            | 12.3 | 17.6        | 30.7        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 344         | 3.9            | 13.3 | 21.5        | 39.0        | 267         | 3.7            | 12.1 | 22.8        | 39.4        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 267         | 4.4            | 5.2  | 72.7        | 92.9        | 204         | 5.3            | 6.3  | 68.9        | 93.3        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 339         | 3.1            | 3.6  | 78.2        | 92.1        | 262         | 2.9            | 3.4  | 77.5        | 90.5        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 705         | 4.2            | 4.9  | 73.7        | 92.5        | 543         | 5.1            | 5.9  | 70.8        | 93.4        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1050        | 5.3            | 6.4  | 68.5        | 92.0        | 747         | 6.8            | 8.2  | 63.1        | 93.1        |
| Female  | 1146        | 3.4            | 3.9  | 78.2        | 93.5        | 831         | 4.1            | 4.8  | 74.9        | 93.5        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 182         | 3.6            | 3.9  | 79.9        | 97.0        | 143         | 3.4            | 3.6  | 80.8        | 98.3        |
| Female  | 157         | 3.4            | 3.8  | 80.1        | 96.0        | 123         | 4.0            | 4.4  | 77.5        | 96.0        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2292        | 3.6            | 17.3 | 14.0        | 29.7        | 1685        | 4.6            | 20.8 | 13.7        | 33.6        |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 2389        | 1.2            | 21.1 | 3.7         | 9.3         | 1747        | 1.6            | 27.2 | 3.2         | 10.4        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 2389        | 0.6            | 13.1 | 3.5         | 6.2         | 1747        | 0.7            | 15.9 | 3.1         | 6.1         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 2389        | 2.6            | 31.8 | 4.0         | 16.1        | 1747        | 3.4            | 35.9 | 4.2         | 19.7        |
| Iodine content in Household Salt (more than 15 ppm) (%) | 939         | 8.5            | 13.4 | 43.2        | 79.5        | 655         | 11.2           | 19.1 | 33.8        | 80.1        |

| Indicators  | Betul       |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 306         | 4.6            | 20.2 | 14.0        | 34.3        | 195         | 6.5            | 27.6 | 12.2        | 41.0        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 311         | 5.7            | 18.6 | 19.7        | 44.8        | 197         | 6.2            | 17.9 | 22.5        | 49.6        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 314         | 3.7            | 16.2 | 15.5        | 31.9        | 199         | 3.5            | 14.4 | 17.5        | 33.2        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 314         | 3.2            | 12.2 | 19.7        | 34.0        | 199         | 4.5            | 17.0 | 17.9        | 38.1        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 314         | 3.0            | 29.9 | 5.0         | 18.8        | 199         | 3.3            | 32.8 | 4.7         | 20.1        |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1116        | 4.7            | 21.7 | 13.0        | 34.0        | 739         | 6.6            | 31.9 | 9.6         | 39.0        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 1116        | 0.5            | 23.8 | 1.2         | 3.3         | 739         | 0.6            | 31.6 | 0.9         | 3.7         |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3143        | 1.4            | 14.4 | 7.0         | 13.4        | 1712        | 1.7            | 25.9 | 3.7         | 11.7        |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 449         | 3.0            | 25.9 | 6.3         | 20.3        | 270         | 2.2            | 27.9 | 4.1         | 14.5        |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3143        | 2.7            | 19.6 | 8.5         | 20.8        | 1712        | 4.0            | 24.6 | 9.2         | 27.5        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 449         | 2.5            | 16.0 | 10.5        | 21.9        | 270         | 4.0            | 21.0 | 11.4        | 29.6        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 162         | 5.4            | 6.3  | 68.8        | 93.3        | 98          | 7.6            | 8.8  | 59.7        | 96.7        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 264         | 3.4            | 3.6  | 79.9        | 98.8        | 164         | 4.6            | 4.9  | 71.7        | 99.0        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 530         | 3.5            | 3.8  | 79.6        | 98.3        | 329         | 5.2            | 5.7  | 69.8        | 98.4        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1207        | 2.5            | 2.6  | 85.6        | 98.6        | 669         | 4.1            | 4.3  | 75.0        | 99.1        |
| Female  | 1283        | 2.9            | 3.1  | 83.5        | 98.2        | 701         | 4.7            | 5.0  | 72.7        | 98.8        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 209         | 2.3            | 2.4  | 85.6        | 99.4        | 133         | 3.4            | 3.5  | 74.9        | 99.6        |
| Female  | 171         | 2.2            | 2.3  | 87.5        | 98.7        | 110         | 3.0            | 3.1  | 81.8        | 99.1        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3561        | 1.1            | 22.8 | 3.0         | 8.2         | 1956        | 1.9            | 45.0 | 1.5         | 10.8        |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3555        | 0.8            | 11.9 | 5.2         | 8.8         | 1956        | 0.8            | 13.6 | 4.4         | 7.9         |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3555        | 0.4            | 24.8 | 0.9         | 2.7         | 1956        | 0.3            | 27.1 | 0.6         | 2.0         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3555        | 2.2            | 24.6 | 5.1         | 14.9        | 1956        | 3.3            | 31.6 | 5.1         | 20.0        |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1317        | 6.4            | 9.1  | 54.5        | 81.7        | 766         | 5.3            | 7.4  | 59.4        | 82.4        |

| Indicators  | Harda       |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 223         | 0.6            | 11.2 | 4.1         | 6.7         | 165         | 0.6            | 11.0 | 4.3         | 7.0         |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 208         | 4.9            | 7.3  | 55.9        | 77.6        | 155         | 5.8            | 8.6  | 52.9        | 78.1        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 226         | 3.2            | 7.3  | 37.5        | 51.9        | 167         | 3.6            | 8.3  | 36.0        | 52.0        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 226         | 1.1            | 35.1 | 1.4         | 6.7         | 167         | 1.2            | 37.9 | 1.4         | 7.4         |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 226         | 2.0            | 21.6 | 5.7         | 14.9        | 167         | 2.1            | 24.0 | 5.2         | 15.0        |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 849         | 3.5            | 9.2  | 30.8        | 46.4        | 593         | 4.1            | 10.5 | 30.4        | 48.5        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 849         | 0.6            | 31.5 | 1.0         | 4.1         | 593         | 0.6            | 42.0 | 0.6         | 3.7         |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 1939        | 2.0            | 13.1 | 11.6        | 20.7        | 1313        | 2.3            | 16.6 | 9.7         | 20.2        |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 355         | 3.3            | 16.1 | 13.7        | 28.7        | 263         | 3.4            | 17.6 | 12.6        | 28.1        |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 1939        | 2.3            | 12.5 | 13.8        | 24.3        | 1313        | 2.7            | 13.4 | 14.5        | 26.7        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 355         | 4.6            | 20.4 | 13.7        | 34.4        | 263         | 5.3            | 22.2 | 14.1        | 37.9        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 179         | 3.6            | 4.2  | 76.7        | 93.2        | 124         | 4.3            | 5.0  | 73.3        | 93.1        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 231         | 5.3            | 5.8  | 70.1        | 97.7        | 165         | 5.8            | 6.6  | 67.7        | 96.9        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 528         | 5.8            | 6.6  | 67.9        | 96.5        | 362         | 6.8            | 7.9  | 63.2        | 95.5        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 947         | 7.6            | 9.2  | 59.5        | 93.3        | 642         | 8.9            | 11.5 | 52.8        | 91.0        |
| Female  | 1043        | 4.6            | 5.1  | 74.1        | 96.4        | 698         | 5.6            | 6.3  | 69.3        | 95.7        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 192         | 13.0           | 16.4 | 40.3        | 95.5        | 139         | 14.2           | 18.9 | 36.2        | 94.1        |
| Female  | 194         | 8.8            | 10.9 | 54.2        | 93.3        | 137         | 9.7            | 12.7 | 49.2        | 91.2        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2226        | 1.4            | 19.4 | 4.7         | 11.0        | 1520        | 1.8            | 24.2 | 4.3         | 12.3        |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 2244        | 1.2            | 21.5 | 3.5         | 9.0         | 1537        | 1.5            | 25.6 | 3.4         | 10.4        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 2244        | 0.8            | 16.1 | 3.4         | 7.0         | 1537        | 0.9            | 18.4 | 3.3         | 7.4         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 2244        | 0.4            | 8.4  | 3.8         | 5.6         | 1537        | 0.4            | 7.6  | 4.0         | 5.6         |
| Iodine content in Household Salt (more than 15 ppm) (%) | 924         | 12.6           | 17.6 | 38.8        | 91.1        | 665         | 15.3           | 22.0 | 31.3        | 91.8        |

| Indicators  | Hoshangabad |                |             |             |             |             |                |             |             |             |
|---|-------------|----------------|-------------|-------------|-------------|-------------|----------------|-------------|-------------|-------------|
|   | Total       |                |             |             |             | Rural       |                |             |             |             |
|   | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |             |             |             |             |                |             |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |             |             |             |             |                |             |             |             |
| Person  | 252         | 6.3            | 20.2        | 18.8        | 46.5        | 191         | 6.6            | 17.7        | 24.1        | 53.3        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |             |             |             |             |                |             |             |             |
| Person  | 285         | 6.7            | 15.1        | 30.4        | 60.0        | 199         | 8.0            | 18.3        | 27.1        | 61.8        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |             |             |             |             |                |             |             |             |
| Person  | 306         | 3.3            | 7.1         | 39.2        | 54.0        | 225         | 2.9            | 6.5         | 37.6        | 50.5        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |             |             |             |             |                |             |             |             |
| Person  | 340         | 5.5            | 11.2        | 36.6        | 60.9        | 249         | 5.6            | 10.7        | 39.6        | 64.4        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |             |             |             |             |                |             |             |             |
| Person  | 340         | 2.5            | 18.6        | 8.8         | 20.4        | 249         | 2.1            | 17.8        | 7.8         | 17.3        |
| <b>Nutritional status 5-18 years</b>                    |             |                |             |             |             |             |                |             |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |             |             |             |             |                |             |             |             |
| Person  | 1261        | 1.7            | 5.3         | 27.9        | 35.4        | 861         | 1.6            | 4.6         | 31.6        | 38.9        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |             |             |             |             |                |             |             |             |
| Person  | 1261        | 0.4            | 24.3        | 0.9         | 2.8         | 861         | 0.4            | 22.1        | 1.0         | 2.8         |
| <b>Nutritional status 18 years and above</b>            |             |                |             |             |             |             |                |             |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person  | 3204        | 3.5            | 36.9        | 4.0         | 20.9        | 1998        | 1.4            | 28.3        | 2.7         | 9.4         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |             |             |             |             |                |             |             |             |
| Person  | 430         | 4.1            | 24.5        | 9.4         | 28.2        | 261         | 4.5            | 27.9        | 8.2         | 28.9        |
| BMI <18.5 for age 18-59 year (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person  | 3204        | 1.7            | 17.4        | 6.5         | 14.3        | 1998        | 2.1            | 16.8        | 8.7         | 18.3        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |             |             |             |             |                |             |             |             |
| Person  | 430         | 4.7            | 26.6        | 9.4         | 30.9        | 261         | 3.7            | 18.7        | 12.5        | 29.3        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |             |             |             |             |                |             |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |             |             |             |             |                |             |             |             |
| Person  | 208         | 2.1            | 2.8         | 72.6        | 82.3        | 147         | 2.8            | 3.7         | 68.3        | 80.9        |
| Anaemia in the age group 5-9 years (%)                  |             |                |             |             |             |             |                |             |             |             |
| Person  | 319         | 1.9            | 2.1         | 85.5        | 94.2        | 217         | 2.4            | 2.6         | 83.6        | 94.8        |
| Anaemia in the age group 10-17 years (%)                |             |                |             |             |             |             |                |             |             |             |
| Person  | 726         | 2.2            | 2.6         | 80.1        | 90.2        | 501         | 1.8            | 2.0         | 83.6        | 91.7        |
| Anaemia in the age group 18-59 years (%)                |             |                |             |             |             |             |                |             |             |             |
| Male  | 1511        | 1.6            | 1.9         | 79.9        | 87.2        | 930         | 1.8            | 2.1         | 80.6        | 88.5        |
| Female  | 1471        | 1.7            | 1.9         | 87.3        | 95.2        | 938         | 2.2            | 2.4         | 85.9        | 96.6        |
| Anaemia in the age group 60 years and above (%)         |             |                |             |             |             |             |                |             |             |             |
| Male  | 199         | 2.9            | 3.3         | 79.4        | 92.8        | 107         | 4.6            | 5.2         | 73.3        | 95.1        |
| Female  | 185         | 2.3            | 2.6         | 83.1        | 93.7        | 126         | 2.2            | 2.4         | 85.0        | 95.0        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |             |             |             |             |                |             |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person  | 3447        | 1.0            | 13.7        | 5.3         | 9.8         | 2075        | 1.1            | 16.5        | 4.6         | 9.7         |
| <b>Hypertension (18 years and above)</b>                |             |                |             |             |             |             |                |             |             |             |
| <b>Above Normal Range</b>                               |             |                |             |             |             |             |                |             |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |             |             |             |             |                |             |             |             |
| Person  | 3615        | 0.7            | 16.4        | 3.0         | 6.4         | 2238        | 0.8            | 20.1        | 2.7         | 6.6         |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |             |             |             |             |                |             |             |             |
| Person  | -           | -              | -           | -           | -           | -           | -              | -           | -           | -           |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |             |             |             |             |                |             |             |             |
| Person  | 3615        | 1.5            | 12.9        | 8.5         | 15.2        | 2238        | 1.8            | 17.2        | 7.1         | 15.4        |
| Iodine content in Household Salt (more than 15 ppm) (%) | <b>1301</b> | <b>12.8</b>    | <b>19.2</b> | <b>35.2</b> | <b>88.0</b> | <b>833</b>  | <b>16.9</b>    | <b>26.2</b> | <b>25.5</b> | <b>90.5</b> |

| Indicators   | Katni       |                |             |             |             |             |                |             |             |             |
|--|-------------|----------------|-------------|-------------|-------------|-------------|----------------|-------------|-------------|-------------|
|  | Total       |                |             |             |             | Rural       |                |             |             |             |
|  | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                        |             |                |             |             |             |             |                |             |             |             |
| Below -2 SD Wasting (Weight for Height) (%)                    |             |                |             |             |             |             |                |             |             |             |
| Person   | 297         | 4.6            | 16.7        | 18.6        | 38.6        | 241         | 4.3            | 22.0        | 11.9        | 31.0        |
| Below -2 SD Stunting (Height for Age) (%)                      |             |                |             |             |             |             |                |             |             |             |
| Person   | 349         | 6.5            | 16.2        | 27.0        | 55.1        | 267         | 8.4            | 19.0        | 27.2        | 62.5        |
| Below -2 SD Underweight (Weight for Age) (%)                   |             |                |             |             |             |             |                |             |             |             |
| Person   | 364         | 5.0            | 9.7         | 40.8        | 62.4        | 285         | 6.6            | 13.6        | 34.4        | 62.5        |
| Below -2 SD Undernourished (BMI for Age) (%)                   |             |                |             |             |             |             |                |             |             |             |
| Person   | 375         | 4.0            | 10.9        | 28.3        | 45.7        | 289         | 4.7            | 18.1        | 16.9        | 37.3        |
| Above 2 SD Overnourished (BMI for Age) (%)                     |             |                |             |             |             |             |                |             |             |             |
| Person   | 375         | 1.0            | 47.5        | 0.8         | 6.1         | 289         | 1.4            | 46.8        | 1.0         | 7.9         |
| <b>Nutritional status 5-18 years</b>                           |             |                |             |             |             |             |                |             |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)                   |             |                |             |             |             |             |                |             |             |             |
| Person   | 1361        | 2.7            | 6.7         | 34.7        | 46.6        | 1031        | 3.2            | 8.0         | 33.6        | 47.7        |
| Above 2 SD Overnourished (BMI for Age) (%)                     |             |                |             |             |             |             |                |             |             |             |
| Person   | 1361        | 0.4            | 36.8        | 0.4         | 2.2         | 1031        | 0.5            | 38.8        | 0.5         | 2.8         |
| <b>Nutritional status 18 years and above</b>                   |             |                |             |             |             |             |                |             |             |             |
| BMI ≥25.0 for age 18-59 year (%)                               |             |                |             |             |             |             |                |             |             |             |
| Person   | 2605        | 1.7            | 18.3        | 6.1         | 13.9        | 1908        | 1.8            | 22.0        | 4.8         | 13.1        |
| BMI ≥25.0 for age 60 year and above (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person   | 355         | 1.8            | 22.3        | 5.0         | 13.4        | 278         | 2.1            | 31.7        | 3.2         | 13.2        |
| BMI <18.5 for age 18-59 year (%)                               |             |                |             |             |             |             |                |             |             |             |
| Person   | 2605        | 1.5            | 5.5         | 24.9        | 31.7        | 1908        | 2.0            | 6.3         | 27.5        | 36.5        |
| BMI <18.5 for age 60 year and above (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person   | 355         | 4.7            | 12.2        | 28.5        | 49.4        | 278         | 5.7            | 12.4        | 33.1        | 58.3        |
| <b>Anaemia status by Haemoglobin Level</b>                     |             |                |             |             |             |             |                |             |             |             |
| Anaemia in the age group 6-59 months (%)                       |             |                |             |             |             |             |                |             |             |             |
| Person   | 273         | 1.7            | 1.8         | 88.4        | 96.0        | 224         | 1.9            | 2.1         | 86.2        | 94.9        |
| Anaemia in the age group 5-9 years (%)                         |             |                |             |             |             |             |                |             |             |             |
| Person   | 439         | 3.8            | 4.2         | 78.5        | 97.3        | 344         | 4.9            | 5.4         | 73.3        | 96.8        |
| Anaemia in the age group 10-17 years (%)                       |             |                |             |             |             |             |                |             |             |             |
| Person   | 739         | 3.7            | 4.4         | 74.8        | 91.4        | 593         | 4.4            | 5.3         | 70.5        | 89.9        |
| Anaemia in the age group 18-59 years (%)                       |             |                |             |             |             |             |                |             |             |             |
| Male   | 1137        | 4.8            | 6.0         | 67.7        | 88.9        | 891         | 6.0            | 7.8         | 60.8        | 86.8        |
| Female   | 1250        | 2.4            | 2.8         | 81.2        | 92.1        | 973         | 2.9            | 3.5         | 77.5        | 90.6        |
| Anaemia in the age group 60 years and above (%)                |             |                |             |             |             |             |                |             |             |             |
| Male   | 147         | 2.9            | 3.1         | 83.7        | 97.8        | 120         | 3.6            | 3.9         | 79.5        | 97.3        |
| Female   | 173         | 2.8            | 3.1         | 82.1        | 95.0        | 145         | 3.5            | 3.9         | 78.8        | 94.8        |
| <b>Blood Sugar Level (18 years and above)</b>                  |             |                |             |             |             |             |                |             |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                               |             |                |             |             |             |             |                |             |             |             |
| Person   | 2798        | 1.1            | 16.3        | 4.7         | 9.7         | 2040        | 1.4            | 18.6        | 4.8         | 10.9        |
| <b>Hypertension (18 years and above)</b>                       |             |                |             |             |             |             |                |             |             |             |
| <b>Above Normal Range</b>                                      |             |                |             |             |             |             |                |             |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)              |             |                |             |             |             |             |                |             |             |             |
| Person   | 2819        | 1.0            | 15.3        | 4.8         | 9.5         | 2052        | 1.5            | 18.8        | 5.1         | 11.7        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)              |             |                |             |             |             |             |                |             |             |             |
| Person   | 2819        | 0.5            | 9.2         | 4.5         | 6.7         | 2052        | 0.6            | 9.1         | 5.2         | 7.7         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)              |             |                |             |             |             |             |                |             |             |             |
| Person   | 2819        | 0.6            | 11.3        | 4.0         | 6.6         | 2052        | 0.7            | 12.5        | 4.5         | 7.7         |
| <b>Iodine content in Household Salt (more than 15 ppm) (%)</b> | <b>1273</b> | <b>7.4</b>     | <b>11.6</b> | <b>46.9</b> | <b>78.6</b> | <b>947</b>  | <b>10.5</b>    | <b>18.5</b> | <b>33.9</b> | <b>77.2</b> |

| Indicators   | Jabalpur    |                |            |             |             |             |                |            |             |             |
|--|-------------|----------------|------------|-------------|-------------|-------------|----------------|------------|-------------|-------------|
|  | Total       |                |            |             |             | Rural       |                |            |             |             |
|  | Sample Size | Standard Error | PRSE       | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE       | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                        |             |                |            |             |             |             |                |            |             |             |
| Below -2 SD Wasting (Weight for Height) (%)                    |             |                |            |             |             |             |                |            |             |             |
| Person   | 224         | 2.3            | 6.7        | 28.9        | 39.0        | 56          | 2.2            | 6.8        | 27.5        | 37.2        |
| Below -2 SD Stunting (Height for Age) (%)                      |             |                |            |             |             |             |                |            |             |             |
| Person   | 219         | 5.5            | 9.8        | 43.3        | 67.3        | 58          | 11.2           | 25.0       | 22.8        | 69.0        |
| Below -2 SD Underweight (Weight for Age) (%)                   |             |                |            |             |             |             |                |            |             |             |
| Person   | 271         | 4.4            | 8.3        | 43.0        | 62.4        | 58          | 2.6            | 5.4        | 42.5        | 54.1        |
| Below -2 SD Undernourished (BMI for Age) (%)                   |             |                |            |             |             |             |                |            |             |             |
| Person   | 271         | 4.1            | 14.1       | 21.0        | 39.3        | 58          | 5.8            | 18.8       | 19.7        | 45.2        |
| Above 2 SD Overnourished (BMI for Age) (%)                     |             |                |            |             |             |             |                |            |             |             |
| Person   | 271         | 7.8            | 27.9       | 14.1        | 47.9        | 58          | 2.2            | 18.5       | 7.9         | 18.0        |
| <b>Nutritional status 5-18 years</b>                           |             |                |            |             |             |             |                |            |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)                   |             |                |            |             |             |             |                |            |             |             |
| Person   | 1446        | 4.1            | 13.4       | 22.2        | 40.3        | 351         | 11.3           | 30.5       | 16.7        | 63.4        |
| Above 2 SD Overnourished (BMI for Age) (%)                     |             |                |            |             |             |             |                |            |             |             |
| Person   | 1446        | 0.5            | 30.2       | 0.8         | 3.1         | -           | -              | -          | -           | -           |
| <b>Nutritional status 18 years and above</b>                   |             |                |            |             |             |             |                |            |             |             |
| BMI ≥25.0 for age 18-59 year (%)                               |             |                |            |             |             |             |                |            |             |             |
| Person   | 3835        | 1.3            | 17.2       | 4.9         | 10.8        | 969         | 0.9            | 18.5       | 3.2         | 7.4         |
| BMI ≥25.0 for age 60 year and above (%)                        |             |                |            |             |             |             |                |            |             |             |
| Person   | 625         | 2.8            | 25.9       | 5.7         | 18.8        | 158         | 1.4            | 37.2       | 1.6         | 8.6         |
| BMI <18.5 for age 18-59 year (%)                               |             |                |            |             |             |             |                |            |             |             |
| Person   | 3835        | 3.0            | 39.5       | 3.1         | 17.8        | -           | -              | -          | -           | -           |
| BMI <18.5 for age 60 year and above (%)                        |             |                |            |             |             |             |                |            |             |             |
| Person   | 625         | 3.8            | 43.2       | 3.2         | 22.0        | -           | -              | -          | -           | -           |
| <b>Anaemia status by Haemoglobin Level</b>                     |             |                |            |             |             |             |                |            |             |             |
| Anaemia in the age group 6-59 months (%)                       |             |                |            |             |             |             |                |            |             |             |
| Person   | 189         | 4.2            | 4.8        | 75.0        | 94.0        | 52          | 4.3            | 4.7        | 75.0        | 98.0        |
| Anaemia in the age group 5-9 years (%)                         |             |                |            |             |             |             |                |            |             |             |
| Person   | 405         | 1.9            | 2.1        | 88.0        | 98.0        | 88          | 1.6            | 1.6        | 90.0        | 100.0       |
| Anaemia in the age group 10-17 years (%)                       |             |                |            |             |             |             |                |            |             |             |
| Person   | 869         | 3.3            | 3.5        | 81.0        | 97.0        | 217         | 3.6            | 3.7        | 69.0        | 100.0       |
| Anaemia in the age group 18-59 years (%)                       |             |                |            |             |             |             |                |            |             |             |
| Male   | 1948        | 1.3            | 1.4        | 92.0        | 98.0        | 513         | 1.8            | 1.8        | 88.0        | 100.0       |
| Female   | 1776        | 2.0            | 2.1        | 88.0        | 98.0        | 458         | 1.9            | 2.0        | 87.0        | 100.0       |
| Anaemia in the age group 60 years and above (%)                |             |                |            |             |             |             |                |            |             |             |
| Male   | 300         | 1.0            | 1.0        | 94.0        | 100.0       | 82          | 1.5            | 1.5        | 85.0        | 100.0       |
| Female   | 316         | 1.7            | 1.7        | 90.0        | 99.0        | 78          | 1.6            | 1.7        | 81.0        | 100.0       |
| <b>Blood Sugar Level (18 years and above)</b>                  |             |                |            |             |             |             |                |            |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                               |             |                |            |             |             |             |                |            |             |             |
| Person   | 4009        | 2.4            | 21.1       | 7.1         | 17.8        | -           | -              | -          | -           | -           |
| <b>Hypertension (18 years and above)</b>                       |             |                |            |             |             |             |                |            |             |             |
| <b>Above Normal Range</b>                                      |             |                |            |             |             |             |                |            |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)              |             |                |            |             |             |             |                |            |             |             |
| Person   | 4414        | 1.5            | 20.6       | 4.6         | 11.3        | -           | -              | -          | -           | -           |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)              |             |                |            |             |             |             |                |            |             |             |
| Person   | 4414        | 0.6            | 15.5       | 2.8         | 5.5         | 1114        | 1.6            | 40.6       | 1.6         | 9.2         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)              |             |                |            |             |             |             |                |            |             |             |
| Person   | 4414        | 1.4            | 23.6       | 3.4         | 9.7         | -           | -              | -          | -           | -           |
| <b>Iodine content in Household Salt (more than 15 ppm) (%)</b> | <b>1513</b> | <b>2.8</b>     | <b>3.7</b> | <b>67.3</b> | <b>79.4</b> | <b>384</b>  | <b>5.1</b>     | <b>7.8</b> | <b>53.7</b> | <b>75.8</b> |



| Indicators  | Narsimhapur |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 261         | 3.7            | 22.2 | 9.9         | 26.8        | 250         | 3.8            | 22.9 | 9.7         | 27.1        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 263         | 4.4            | 15.7 | 19.1        | 38.8        | 253         | 4.5            | 15.6 | 19.7        | 39.7        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 270         | 3.3            | 12.6 | 19.7        | 34.8        | 259         | 3.4            | 12.7 | 20.1        | 35.6        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 270         | 3.0            | 15.2 | 13.8        | 27.4        | 259         | 3.1            | 15.6 | 13.6        | 27.4        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | -           | -              | -    | -           | -           | -           | -              | -    | -           | -           |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1009        | 3.7            | 7.6  | 39.9        | 56.3        | 959         | 3.7            | 7.8  | 39.8        | 56.6        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 1009        | 0.2            | 41.5 | 0.2         | 1.5         | 959         | 0.3            | 48.2 | 0.2         | 1.6         |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2088        | 2.8            | 26.9 | 5.7         | 18.8        | 1940        | 3.0            | 29.7 | 5.0         | 19.1        |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 411         | 3.0            | 26.3 | 6.3         | 20.2        | 377         | 3.1            | 29.0 | 5.4         | 20.0        |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2088        | 4.1            | 12.3 | 24.7        | 43.1        | 1940        | 4.2            | 12.6 | 24.8        | 43.7        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 411         | 6.6            | 17.7 | 23.9        | 53.0        | 377         | 6.8            | 17.9 | 24.2        | 54.1        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 176         | 5.8            | 8.2  | 56.5        | 82.4        | 169         | 6.0            | 8.4  | 56.1        | 82.6        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 242         | 1.6            | 1.6  | 90.5        | 98.2        | 229         | 1.5            | 1.6  | 90.7        | 98.7        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 520         | 2.2            | 2.3  | 86.0        | 96.2        | 493         | 2.3            | 2.4  | 86.2        | 97.1        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 950         | 2.7            | 3.0  | 81.9        | 94.7        | 877         | 2.8            | 3.1  | 82.2        | 95.8        |
| Female  | 918         | 1.9            | 2.1  | 87.9        | 97.3        | 843         | 2.1            | 2.2  | 87.4        | 97.6        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 176         | 4.5            | 4.8  | 72.9        | 98.8        | 157         | 4.7            | 5.0  | 70.6        | 99.0        |
| Female  | 167         | 4.9            | 5.5  | 72.2        | 96.9        | 150         | 5.2            | 5.8  | 71.0        | 97.1        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2466        | 1.5            | 12.4 | 9.1         | 15.9        | 2287        | 1.5            | 12.9 | 8.9         | 15.9        |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 2466        | 0.3            | 20.8 | 1.0         | 2.6         | 2288        | 0.4            | 23.7 | 0.9         | 2.6         |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 2466        | 1.6            | 21.3 | 4.7         | 12.4        | 2288        | 1.7            | 22.9 | 4.4         | 12.4        |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | -           | -              | -    | -           | -           | -           | -              | -    | -           | -           |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1097        | 8.4            | 11.6 | 50.3        | 87.1        | 1019        | 8.7            | 12.1 | 48.9        | 86.8        |

| Indicators  | Dindori     |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 455         | 3.3            | 12.9 | 19.2        | 34.0        | 408         | 3.4            | 13.4 | 18.7        | 33.9        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 403         | 4.0            | 6.2  | 55.4        | 73.2        | 361         | 4.2            | 6.4  | 55.3        | 73.6        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 476         | 2.8            | 4.8  | 51.4        | 63.7        | 427         | 2.8            | 4.9  | 51.5        | 64.2        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 476         | 3.3            | 14.8 | 15.8        | 30.4        | 427         | 3.4            | 15.2 | 15.5        | 30.4        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 476         | 1.6            | 7.1  | 19.0        | 26.1        | 427         | 1.6            | 7.3  | 18.9        | 26.1        |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1620        | 3.9            | 11.1 | 27.1        | 44.3        | 1478        | 4.0            | 11.2 | 27.2        | 44.8        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 1620        | 0.3            | 12.6 | 1.6         | 2.8         | 1478        | 0.3            | 12.3 | 1.6         | 2.7         |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3186        | 1.8            | 27.0 | 3.6         | 12.1        | 2825        | 1.9            | 28.6 | 3.4         | 12.4        |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 479         | 1.4            | 18.2 | 5.0         | 11.4        | 432         | 1.4            | 18.6 | 5.0         | 11.4        |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3186        | 1.3            | 5.0  | 22.3        | 28.2        | 2825        | 1.3            | 5.3  | 22.7        | 28.6        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 479         | 3.3            | 10.0 | 26.5        | 41.3        | 432         | 3.4            | 10.2 | 26.4        | 41.7        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 363         | 3.4            | 3.7  | 79.8        | 95.5        | 332         | 3.4            | 3.8  | 79.8        | 95.9        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 567         | 1.3            | 1.4  | 91.3        | 97.7        | 532         | 1.4            | 1.4  | 91.3        | 97.8        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 855         | 2.0            | 2.1  | 87.5        | 97.2        | 765         | 2.1            | 2.2  | 87.4        | 97.5        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1465        | 3.0            | 3.3  | 81.8        | 96.1        | 1293        | 3.1            | 3.3  | 81.9        | 96.6        |
| Female  | 1558        | 2.5            | 2.7  | 85.1        | 97.6        | 1373        | 2.6            | 2.8  | 84.8        | 97.8        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 227         | 2.1            | 2.2  | 87.5        | 97.4        | 203         | 2.1            | 2.2  | 87.7        | 97.9        |
| Female  | 244         | 2.3            | 2.5  | 86.3        | 97.9        | 218         | 2.4            | 2.6  | 85.8        | 97.9        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3559        | 1.0            | 18.4 | 3.5         | 7.9         | 3172        | 1.0            | 20.3 | 3.2         | 7.8         |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3577        | 0.7            | 11.4 | 4.9         | 8.1         | 3186        | 0.7            | 11.7 | 4.9         | 8.2         |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3577        | 0.4            | 11.0 | 2.9         | 4.7         | 3186        | 0.4            | 11.2 | 2.9         | 4.7         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3577        | 5.5            | 42.4 | 4.8         | 30.5        | 3186        | 5.7            | 43.3 | 4.7         | 31.3        |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1447        | 11.5           | 18.5 | 35.5        | 82.8        | 1293        | 11.9           | 19.5 | 33.9        | 82.7        |

| Indicators  | Mandla      |                |             |             |             |             |                |             |             |             |
|---|-------------|----------------|-------------|-------------|-------------|-------------|----------------|-------------|-------------|-------------|
|   | Total       |                |             |             |             | Rural       |                |             |             |             |
|   | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE        | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |             |             |             |             |                |             |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |             |             |             |             |                |             |             |             |
| Person  | 303         | 2.2            | 7.9         | 22.7        | 32.2        | 260         | 2.6            | 9.7         | 21.2        | 32.7        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |             |             |             |             |                |             |             |             |
| Person  | 310         | 1.7            | 3.9         | 39.2        | 46.6        | 265         | 2.0            | 4.6         | 38.2        | 47.0        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |             |             |             |             |                |             |             |             |
| Person  | 317         | 2.8            | 6.0         | 40.7        | 53.2        | 271         | 3.2            | 7.1         | 38.7        | 53.0        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |             |             |             |             |                |             |             |             |
| Person  | 318         | 2.7            | 10.2        | 21.0        | 33.0        | 272         | 2.9            | 11.9        | 18.8        | 31.9        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |             |             |             |             |                |             |             |             |
| Person  | -           | -              | -           | -           | -           | -           | -              | -           | -           | -           |
| <b>Nutritional status 5-18 years</b>                    |             |                |             |             |             |             |                |             |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |             |             |             |             |                |             |             |             |
| Person  | 1140        | 3.1            | 8.7         | 28.7        | 42.3        | 1010        | 3.4            | 10.0        | 26.8        | 41.7        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |             |             |             |             |                |             |             |             |
| Person  | 1140        | 0.3            | 21.0        | 1.0         | 2.6         | 1010        | 0.3            | 24.3        | 0.8         | 2.4         |
| <b>Nutritional status 18 years and above</b>            |             |                |             |             |             |             |                |             |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person  | 2379        | 1.8            | 29.7        | 3.1         | 11.7        | 2069        | 0.6            | 14.6        | 2.9         | 5.7         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |             |             |             |             |                |             |             |             |
| Person  | 317         | 2.3            | 32.7        | 3.3         | 14.3        | 256         | 1.9            | 40.2        | 1.9         | 11.4        |
| BMI <18.5 for age 18-59 year (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person  | 2379        | 2.1            | 6.4         | 27.9        | 37.4        | 2069        | 1.9            | 5.4         | 30.4        | 39.0        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |             |             |             |             |                |             |             |             |
| Person  | 317         | 4.7            | 11.3        | 31.8        | 52.7        | 256         | 4.5            | 9.5         | 37.1        | 57.2        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |             |             |             |             |                |             |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |             |             |             |             |                |             |             |             |
| Person  | 206         | 2.3            | 2.7         | 79.3        | 89.5        | 182         | 2.2            | 2.7         | 78.5        | 88.5        |
| Anaemia in the age group 5-9 years (%)                  |             |                |             |             |             |             |                |             |             |             |
| Person  | 366         | 1.2            | 1.2         | 92.6        | 98.3        | 330         | 1.3            | 1.4         | 92.0        | 98.4        |
| Anaemia in the age group 10-17 years (%)                |             |                |             |             |             |             |                |             |             |             |
| Person  | 635         | 3.5            | 4.0         | 78.2        | 94.6        | 559         | 4.1            | 4.6         | 75.8        | 95.0        |
| Anaemia in the age group 18-59 years (%)                |             |                |             |             |             |             |                |             |             |             |
| Male  | 1052        | 4.0            | 4.7         | 74.7        | 93.1        | 924         | 4.6            | 5.4         | 72.1        | 93.3        |
| Female  | 1233        | 2.4            | 2.7         | 84.0        | 95.3        | 1074        | 2.7            | 2.9         | 82.4        | 94.6        |
| Anaemia in the age group 60 years and above (%)         |             |                |             |             |             |             |                |             |             |             |
| Male  | 129         | 4.5            | 4.9         | 75.0        | 96.9        | 107         | 5.5            | 6.2         | 69.6        | 96.9        |
| Female  | 172         | 1.7            | 1.8         | 88.9        | 97.1        | 137         | 1.9            | 2.0         | 88.3        | 97.3        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |             |             |             |             |                |             |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |             |             |             |             |                |             |             |             |
| Person  | 2623        | 2.0            | 18.3        | 7.2         | 16.3        | 2256        | 2.1            | 22.1        | 5.6         | 15.0        |
| <b>Hypertension (18 years and above)</b>                |             |                |             |             |             |             |                |             |             |             |
| <b>Above Normal Range</b>                               |             |                |             |             |             |             |                |             |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |             |             |             |             |                |             |             |             |
| Person  | 2633        | 1.4            | 15.2        | 6.4         | 12.5        | 2264        | 1.6            | 18.1        | 5.8         | 13.0        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |             |             |             |             |                |             |             |             |
| Person  | 2633        | 0.7            | 9.8         | 5.4         | 8.3         | 2264        | 0.7            | 11.4        | 5.1         | 8.4         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |             |             |             |             |                |             |             |             |
| Person  | 2633        | 0.8            | 13.1        | 4.3         | 7.7         | 2264        | 0.8            | 13.5        | 4.4         | 8.0         |
| Iodine content in Household Salt (more than 15 ppm) (%) | <b>1177</b> | <b>8.0</b>     | <b>12.9</b> | <b>43.5</b> | <b>78.0</b> | <b>1020</b> | <b>8.9</b>     | <b>14.6</b> | <b>40.4</b> | <b>78.1</b> |

| Indicators  | Chindwara   |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 263         | 3.1            | 14.8 | 14.8        | 28.8        | 186         | 3.8            | 19.4 | 12.4        | 29.7        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 271         | 3.9            | 11.1 | 26.9        | 44.3        | 197         | 4.8            | 13.4 | 26.0        | 47.4        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 284         | 3.6            | 10.2 | 27.3        | 43.2        | 204         | 4.5            | 11.6 | 29.0        | 48.9        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 284         | 3.0            | 13.1 | 16.7        | 30.0        | 205         | 3.4            | 14.5 | 16.6        | 32.0        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 284         | 1.5            | 15.0 | 7.0         | 13.8        | 205         | 1.2            | 10.8 | 8.7         | 14.1        |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1006        | 0.7            | 7.8  | 7.0         | 10.0        | 741         | 0.6            | 7.0  | 6.8         | 9.3         |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 1006        | 0.4            | 49.9 | 0.2         | 2.3         | -           | -              | -    | -           | -           |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2971        | 0.7            | 20.0 | 2.1         | 5.4         | 2184        | 0.8            | 26.3 | 1.8         | 5.6         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 307         | 1.2            | 21.5 | 3.4         | 9.0         | 228         | 1.4            | 25.0 | 3.2         | 9.8         |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2971        | 0.4            | 19.6 | 1.4         | 3.2         | 2184        | 0.4            | 30.3 | 0.8         | 2.7         |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 307         | 0.7            | 33.8 | 1.0         | 4.4         | 228         | 0.7            | 46.1 | 0.6         | 4.3         |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 180         | 5.0            | 6.8  | 61.6        | 84.2        | 132         | 5.2            | 6.7  | 63.9        | 87.3        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 291         | 2.6            | 2.9  | 83.7        | 96.3        | 220         | 3.1            | 3.3  | 81.3        | 98.8        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 639         | 2.8            | 3.0  | 83.2        | 97.2        | 489         | 2.9            | 3.0  | 82.9        | 98.5        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1498        | 1.1            | 1.1  | 93.3        | 98.5        | 1130        | 0.9            | 0.9  | 94.8        | 99.5        |
| Female  | 1399        | 3.4            | 3.7  | 79.4        | 95.5        | 1031        | 2.5            | 2.7  | 85.2        | 97.7        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 154         | 2.3            | 2.4  | 86.4        | 98.7        | 109         | 2.6            | 2.7  | 82.2        | 99.4        |
| Female  | 155         | 5.0            | 5.6  | 71.9        | 96.1        | 123         | 4.1            | 4.4  | 76.7        | 97.6        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3248        | 0.6            | 9.0  | 5.6         | 8.5         | 2396        | 0.7            | 9.9  | 5.8         | 9.0         |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3249        | 0.7            | 17.0 | 3.0         | 6.4         | 2399        | 0.8            | 17.1 | 3.2         | 6.9         |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3249        | 0.6            | 23.1 | 1.6         | 4.6         | 2399        | 0.8            | 24.3 | 1.9         | 5.5         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3249        | 1.2            | 10.7 | 8.8         | 14.3        | 2399        | 1.5            | 13.1 | 8.5         | 15.4        |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1208        | 2.6            | 3.6  | 67.0        | 78.9        | 890         | 3.2            | 4.3  | 65.3        | 79.6        |

| Indicators  | Seoni       |                |            |             |             |             |                |            |             |             |
|---|-------------|----------------|------------|-------------|-------------|-------------|----------------|------------|-------------|-------------|
|   | Total       |                |            |             |             | Rural       |                |            |             |             |
|   | Sample Size | Standard Error | PRSE       | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE       | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |            |             |             |             |                |            |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |            |             |             |             |                |            |             |             |
| Person  | 242         | 2.6            | 17.8       | 9.6         | 21.5        | 187         | 2.0            | 18.6       | 6.9         | 16.0        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |            |             |             |             |                |            |             |             |
| Person  | 209         | 6.8            | 11.6       | 42.6        | 72.2        | 166         | 7.5            | 12.6       | 42.1        | 74.8        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |            |             |             |             |                |            |             |             |
| Person  | 274         | 4.4            | 8.0        | 44.8        | 64.4        | 216         | 5.4            | 9.8        | 42.4        | 66.2        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |            |             |             |             |                |            |             |             |
| Person  | 276         | 2.0            | 18.8       | 6.9         | 16.0        | 218         | 1.6            | 21.4       | 4.5         | 11.9        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |            |             |             |             |                |            |             |             |
| Person  | 276         | 4.0            | 17.4       | 15.1        | 33.0        | 218         | 4.4            | 18.3       | 15.5        | 35.2        |
| <b>Nutritional status 5-18 years</b>                    |             |                |            |             |             |             |                |            |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |            |             |             |             |                |            |             |             |
| Person  | 1092        | 3.9            | 13.0       | 22.2        | 39.5        | 1007        | 4.1            | 13.5       | 22.3        | 40.6        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |            |             |             |             |                |            |             |             |
| Person  | 1092        | 0.8            | 29.4       | 1.3         | 4.9         | 1007        | 0.8            | 31.9       | 1.2         | 4.8         |
| <b>Nutritional status 18 years and above</b>            |             |                |            |             |             |             |                |            |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |            |             |             |             |                |            |             |             |
| Person  | 2587        | 0.6            | 9.0        | 5.0         | 7.7         | 2223        | 0.4            | 8.8        | 3.6         | 5.4         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |            |             |             |             |                |            |             |             |
| Person  | 315         | 1.3            | 16.7       | 5.3         | 11.3        | 273         | 0.9            | 14.7       | 4.6         | 8.8         |
| BMI <18.5 for age 18-59 year (%)                        |             |                |            |             |             |             |                |            |             |             |
| Person  | 2587        | 1.2            | 5.8        | 18.8        | 24.2        | 2223        | 1.5            | 6.6        | 18.9        | 25.7        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |            |             |             |             |                |            |             |             |
| Person  | 315         | 3.5            | 9.7        | 28.5        | 44.2        | 273         | 3.9            | 10.1       | 29.7        | 47.2        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |            |             |             |             |                |            |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |            |             |             |             |                |            |             |             |
| Person  | 213         | 0.9            | 1.0        | 94.3        | 99.3        | 183         | 0.8            | 0.8        | 94.2        | 99.8        |
| Anaemia in the age group 5-9 years (%)                  |             |                |            |             |             |             |                |            |             |             |
| Person  | 318         | 1.3            | 1.3        | 92.0        | 99.3        | 294         | 1.4            | 1.4        | 91.5        | 99.4        |
| Anaemia in the age group 10-17 years (%)                |             |                |            |             |             |             |                |            |             |             |
| Person  | 651         | 2.9            | 3.2        | 82.8        | 96.8        | 596         | 3.0            | 3.2        | 82.6        | 97.7        |
| Anaemia in the age group 18-59 years (%)                |             |                |            |             |             |             |                |            |             |             |
| Male  | 1269        | 2.7            | 3.0        | 83.0        | 95.7        | 1096        | 2.8            | 3.0        | 83.5        | 97.4        |
| Female  | 1254        | 1.8            | 1.8        | 89.5        | 98.1        | 1073        | 1.7            | 1.8        | 89.7        | 98.5        |
| Anaemia in the age group 60 years and above (%)         |             |                |            |             |             |             |                |            |             |             |
| Male  | 161         | 2.7            | 2.9        | 84.1        | 98.6        | 139         | 2.5            | 2.6        | 84.9        | 99.1        |
| Female  | 143         | 1.2            | 1.2        | 92.3        | 99.6        | 126         | 1.2            | 1.2        | 91.2        | 99.8        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |            |             |             |             |                |            |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |            |             |             |             |                |            |             |             |
| Person  | 2857        | 0.8            | 12.5       | 4.6         | 8.1         | 2465        | 0.7            | 12.3       | 4.3         | 7.4         |
| <b>Hypertension (18 years and above)</b>                |             |                |            |             |             |             |                |            |             |             |
| <b>Above Normal Range</b>                               |             |                |            |             |             |             |                |            |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |            |             |             |             |                |            |             |             |
| Person  | 2864        | 0.6            | 11.0       | 4.2         | 6.9         | 2470        | 0.7            | 12.8       | 3.9         | 6.9         |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |            |             |             |             |                |            |             |             |
| Person  | 2864        | 0.9            | 19.2       | 3.1         | 7.2         | 2470        | 1.0            | 21.2       | 3.0         | 7.6         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |            |             |             |             |                |            |             |             |
| Person  | 2864        | 1.8            | 28.7       | 3.2         | 11.5        | 2470        | 2.0            | 31.8       | 3.1         | 12.5        |
| Iodine content in Household Salt (more than 15 ppm) (%) | <b>1106</b> | <b>4.3</b>     | <b>5.1</b> | <b>72.8</b> | <b>92.3</b> | <b>963</b>  | <b>4.8</b>     | <b>5.6</b> | <b>71.1</b> | <b>92.9</b> |

| Indicators  | Balaghat    |                |      |             |             |             |                |      |             |             |
|---|-------------|----------------|------|-------------|-------------|-------------|----------------|------|-------------|-------------|
|   | Total       |                |      |             |             | Rural       |                |      |             |             |
|   | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit | Sample Size | Standard Error | PRSE | Lower Limit | Upper Limit |
| <b>Nutritional status below 5 years</b>                 |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Wasting (Weight for Height) (%)             |             |                |      |             |             |             |                |      |             |             |
| Person  | 332         | 4.4            | 19.6 | 14.5        | 33.6        | 300         | 4.9            | 20.0 | 15.5        | 36.6        |
| Below -2 SD Stunting (Height for Age) (%)               |             |                |      |             |             |             |                |      |             |             |
| Person  | 335         | 4.9            | 14.7 | 23.7        | 44.7        | 303         | 5.5            | 16.9 | 21.8        | 45.1        |
| Below -2 SD Underweight (Weight for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 342         | 4.4            | 12.7 | 25.8        | 44.7        | 310         | 4.8            | 13.5 | 26.3        | 46.9        |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 341         | 5.5            | 25.3 | 12.2        | 36.1        | 308         | 6.1            | 24.6 | 13.9        | 39.9        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 341         | 1.3            | 30.9 | 2.2         | 8.2         | 308         | 1.5            | 37.2 | 1.8         | 8.8         |
| <b>Nutritional status 5-18 years</b>                    |             |                |      |             |             |             |                |      |             |             |
| Below -2 SD Undernourished (BMI for Age) (%)            |             |                |      |             |             |             |                |      |             |             |
| Person  | 1059        | 2.4            | 6.3  | 32.4        | 42.7        | 911         | 2.2            | 5.8  | 32.8        | 42.4        |
| Above 2 SD Overnourished (BMI for Age) (%)              |             |                |      |             |             |             |                |      |             |             |
| Person  | 1059        | 0.4            | 32.2 | 0.7         | 2.8         | 911         | 0.5            | 41.3 | 0.5         | 3.0         |
| <b>Nutritional status 18 years and above</b>            |             |                |      |             |             |             |                |      |             |             |
| BMI ≥25.0 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2778        | 1.0            | 16.5 | 4.1         | 8.7         | 2417        | 0.8            | 17.7 | 3.1         | 6.8         |
| BMI ≥25.0 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 394         | 2.1            | 26.4 | 4.4         | 14.2        | 320         | 2.6            | 41.4 | 2.3         | 15.4        |
| BMI <18.5 for age 18-59 year (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 2778        | 3.1            | 7.7  | 32.8        | 46.8        | 2417        | 3.4            | 8.4  | 33.3        | 48.6        |
| BMI <18.5 for age 60 year and above (%)                 |             |                |      |             |             |             |                |      |             |             |
| Person  | 394         | 3.9            | 9.0  | 34.4        | 51.9        | 320         | 4.7            | 10.8 | 33.6        | 54.6        |
| <b>Anaemia status by Haemoglobin Level</b>              |             |                |      |             |             |             |                |      |             |             |
| Anaemia in the age group 6-59 months (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 251         | 2.7            | 3.2  | 79.6        | 91.6        | 229         | 2.9            | 3.4  | 78.0        | 90.8        |
| Anaemia in the age group 5-9 years (%)                  |             |                |      |             |             |             |                |      |             |             |
| Person  | 332         | 2.9            | 3.0  | 83.5        | 97.9        | 282         | 3.3            | 3.5  | 81.3        | 97.9        |
| Anaemia in the age group 10-17 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Person  | 558         | 3.5            | 3.9  | 79.9        | 97.0        | 484         | 4.2            | 4.7  | 76.4        | 96.7        |
| Anaemia in the age group 18-59 years (%)                |             |                |      |             |             |             |                |      |             |             |
| Male  | 1189        | 5.3            | 6.0  | 71.4        | 95.2        | 1033        | 6.3            | 7.3  | 66.8        | 95.6        |
| Female  | 1432        | 2.6            | 2.7  | 85.5        | 97.8        | 1249        | 3.0            | 3.2  | 83.1        | 97.4        |
| Anaemia in the age group 60 years and above (%)         |             |                |      |             |             |             |                |      |             |             |
| Male  | 185         | 5.8            | 6.3  | 68.6        | 98.2        | 153         | 7.2            | 8.0  | 61.9        | 98.0        |
| Female  | 182         | 2.0            | 2.1  | 88.1        | 97.2        | 151         | 2.3            | 2.5  | 86.4        | 96.9        |
| <b>Blood Sugar Level (18 years and above)</b>           |             |                |      |             |             |             |                |      |             |             |
| Blood Sugar Level ≥110 mg/dl (%)                        |             |                |      |             |             |             |                |      |             |             |
| Person  | 3146        | 1.4            | 17.8 | 5.2         | 11.1        | 2720        | 1.5            | 22.8 | 4.1         | 10.9        |
| <b>Hypertension (18 years and above)</b>                |             |                |      |             |             |             |                |      |             |             |
| <b>Above Normal Range</b>                               |             |                |      |             |             |             |                |      |             |             |
| Systolic ≥140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3099        | 1.2            | 15.9 | 5.4         | 10.7        | 2669        | 1.4            | 20.2 | 4.3         | 10.3        |
| Systolic ≥140mm of Hg & Diastolic <90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3099        | 0.4            | 7.2  | 4.2         | 5.7         | 2669        | 0.4            | 8.6  | 3.9         | 5.6         |
| Systolic <140mm of Hg & Diastolic ≥90mm of Hg (%)       |             |                |      |             |             |             |                |      |             |             |
| Person  | 3099        | 0.9            | 14.2 | 4.5         | 8.2         | 2669        | 1.0            | 15.8 | 4.4         | 8.6         |
| Iodine content in Household Salt (more than 15 ppm) (%) | 1250        | 7.4            | 10.0 | 55.7        | 86.8        | 1082        | 8.6            | 12.2 | 49.4        | 85.0        |

Note: Confidence Intervals are computed by logit transformation method using SPSS application.